

# Computing the delta set of an affine semigroup: a status report

Christopher O'Neill

San Diego State University

*cdoneill@sdsu.edu*

Joint with Thomas Barron\* and Roberto Pelayo  
Joint with Pedro García Sánchez and Gautam Webb\*

\* = undergraduate student

Feb 10, 2020

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}.$$

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}$ . “McNugget Semigroup”

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}$ . “McNugget Semigroup”

Factorizations:

$$60 =$$

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}$ . “McNugget Semigroup”

Factorizations:

$$60 = 7(6) + 2(9)$$

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}$ . “McNugget Semigroup”

Factorizations:

$$\begin{aligned} 60 &= 7(6) + 2(9) \\ &= \qquad\qquad\qquad 3(20) \end{aligned}$$

## Definition

A *numerical semigroup*  $S \subset \mathbb{Z}_{\geq 0}$ : closed under **addition**.

## Example

$McN = \langle 6, 9, 20 \rangle = \{0, 6, 9, 12, 15, 18, 20, \dots\}$ . “McNugget Semigroup”

Factorizations:

$$\begin{array}{rcll} 60 & = & 7(6) + 2(9) & \rightsquigarrow (7, 2, 0) \\ & = & 3(20) & \rightsquigarrow (0, 0, 3) \end{array}$$



# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$


# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$




# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$




# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$



$$Z((6, 6)) = \left\{ \begin{array}{l} (6, 0, 0, 0, 0, 0, 0), \\ (0, 1, 0, 0, 1, 0, 0) \end{array} \right\}$$


# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$



$$Z((6, 6)) = \left\{ \begin{array}{l} (6, 0, 0, 0, 0, 0, 0), \\ (0, 1, 0, 0, 1, 0, 0) \end{array} \right\}$$

$$\begin{aligned} (6, 6) &= 6(1, 1) \\ &= (1, 5) + (5, 1) \end{aligned}$$


# Affine semigroups

## Definition

An *affine semigroup*  $S \subset \mathbb{Z}_{\geq 0}^d$ : closed under **vector addition**.

## Example

$$S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$$



$$Z((6, 6)) = \left\{ \begin{array}{l} (6, 0, 0, 0, 0, 0, 0), \\ (0, 1, 0, 0, 1, 0, 0) \end{array} \right\}$$

$$\begin{aligned} (6, 6) &= 6(1, 1) \\ &= (1, 5) + (5, 1) \end{aligned}$$

$S$  numerical semigroup  $\iff d = 1$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .



# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Possible factorization lengths for  $n = 60$ : 3, 7, 8, 9, 10.

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Possible factorization lengths for  $n = 60$ : 3, 7, 8, 9, 10.

$$Z(1000001) =$$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Possible factorization lengths for  $n = 60$ : 3, 7, 8, 9, 10.

$$Z(1000001) = \left\{ \underbrace{\hspace{10em}}_{\text{shortest}}, \dots, \underbrace{\hspace{10em}}_{\text{longest}} \right\}$$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Possible factorization lengths for  $n = 60$ : 3, 7, 8, 9, 10.

$$Z(1000001) = \left\{ \underbrace{(2, 1, 49999)}_{\text{shortest}}, \dots, \underbrace{\hspace{10em}}_{\text{longest}} \right\}$$

# Affine semigroups

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the *set of factorizations* of  $n \in S$ .

$$|\mathbf{a}| = a_1 + \dots + a_k \text{ (length of } \mathbf{a} \text{)}$$

## Example

$S = \langle 6, 9, 20 \rangle$ :

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Possible factorization lengths for  $n = 60$ : 3, 7, 8, 9, 10.

$$Z(1000001) = \left\{ \underbrace{(2, 1, 49999)}_{\text{shortest}}, \dots, \underbrace{(166662, 1, 1)}_{\text{longest}} \right\}$$

# Affine semigroups (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$



# Affine semigroups (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

A geometric viewpoint: non-negative integer solutions to linear equations.

# Affine semigroups (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

A geometric viewpoint: non-negative integer solutions to linear equations.

Example:  $S = \langle 6, 9, 20 \rangle$ ,  $n = 60$ .


# Affine semigroups (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

A geometric viewpoint: non-negative integer solutions to linear equations.

Example:  $S = \langle 6, 9, 20 \rangle$ ,  $n = 60$ .



$$Z(60) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : 60 = 6a_1 + 9a_2 + 20a_3 \right\}$$


# Affine semigroups (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

A geometric viewpoint: non-negative integer solutions to linear equations.

Example:  $S = \langle 6, 9, 20 \rangle$ ,  $n = 60$ .



# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$\begin{aligned} L(n) &= \{ |\mathbf{a}| : \mathbf{a} \in Z(n) \} \\ &= \{ l_1 < \dots < l_r \} \end{aligned}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$\begin{aligned} L(n) &= \{ |\mathbf{a}| : \mathbf{a} \in Z(n) \} \\ &= \{ l_1 < \dots < l_r \} \end{aligned}$$

## Definition

The delta set of  $n \in S$  is  $\Delta(n) = \{ l_2 - l_1, \dots, l_r - l_{r-1} \}$



# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$\begin{aligned} L(n) &= \{ |\mathbf{a}| : \mathbf{a} \in Z(n) \} \\ &= \{ l_1 < \dots < l_r \} \end{aligned}$$

## Definition

The delta set of  $n \in S$  is  $\Delta(n) = \{ l_2 - l_1, \dots, l_r - l_{r-1} \}$

## Example

$$S = \langle 6, 9, 20 \rangle : Z(60) = \{ (10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3) \}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$\begin{aligned} L(n) &= \{ |\mathbf{a}| : \mathbf{a} \in Z(n) \} \\ &= \{ \ell_1 < \dots < \ell_r \} \end{aligned}$$

## Definition

The delta set of  $n \in S$  is  $\Delta(n) = \{ \ell_2 - \ell_1, \dots, \ell_r - \ell_{r-1} \}$

## Example

$$S = \langle 6, 9, 20 \rangle: Z(60) = \{ (10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3) \}$$

$$L(60) = \{ 3, 7, 8, 9, 10 \}$$

$$\Delta(60) = \{ 1, 4 \}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$\begin{aligned} L(n) &= \{ |\mathbf{a}| : \mathbf{a} \in Z(n) \} \\ &= \{ \ell_1 < \dots < \ell_r \} \end{aligned}$$

## Definition

The delta set of  $n \in S$  is  $\Delta(n) = \{ \ell_2 - \ell_1, \dots, \ell_r - \ell_{r-1} \}$

## Example

$$S = \langle 6, 9, 20 \rangle: Z(60) = \{ (10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3) \}$$

$$L(60) = \{ 3, 7, 8, 9, 10 \}$$

$$\Delta(60) = \{ 1, 4 \}$$

$$L(142) = \{ 10, 11, 12, 14, 15, 16, 17, 18, 19 \}$$

$$\Delta(142) = \{ 1, 2 \}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Facts for large $n \in S$

$\min \Delta(n)$  is as small as possible for  $S$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Facts for large $n \in S$

$\min \Delta(n)$  is as small as possible for  $S$

$S = \langle 3, 5, 7 \rangle$ :

$$L(110) = \{16, 18, \dots, 34, 36\}, \quad \Delta(110) = \{2\}$$

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Facts for large $n \in S$

$\min \Delta(n)$  is as small as possible for  $S$

$S = \langle 3, 5, 7 \rangle$ :

$$L(110) = \{16, 18, \dots, 34, 36\}, \quad \Delta(110) = \{2\}$$

$L(n)$  is an arithmetic sequence with a few values removed near the ends

# Delta set

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Facts for large $n \in S$

$\min \Delta(n)$  is as small as possible for  $S$

$S = \langle 3, 5, 7 \rangle$ :

$$L(110) = \{16, 18, \dots, 34, 36\}, \quad \Delta(110) = \{2\}$$

$L(n)$  is an arithmetic sequence with a few values removed near the ends

$S = \langle 42, 86, 245, 285, 365, 463 \rangle$ :

$$L(3023) = \{7, 9, 11, 12, \dots, 46, 47, 58, 62, 64\}, \quad \Delta(3023) = \{1, 2, 4, 9\}$$



# Delta set (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Delta set (a geometric viewpoint)

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

A geometric viewpoint: lattice width

# Delta set (a geometric viewpoint)


Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

A geometric viewpoint: lattice width



# Delta set (a geometric viewpoint)


Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

A geometric viewpoint: lattice width



$$|\mathbf{a}| = \|\mathbf{a}\|_1 \quad (\text{the } \ell_1\text{-norm})$$

# Delta set (a geometric viewpoint)


Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

A geometric viewpoint: lattice width



$$|\mathbf{a}| = \|\mathbf{a}\|_1 \quad (\text{the } \ell_1\text{-norm})$$

$$\min \Delta(n): \min \ell_1 \text{ width}$$

# Delta set (a geometric viewpoint)


Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

A geometric viewpoint: lattice width



$|\mathbf{a}| = \|\mathbf{a}\|_1$  (the  $\ell_1$ -norm)

$\min \Delta(n)$ : min  $\ell_1$  width

extremal lengths near vertices

# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Definition

The delta set of  $S$  is the union  $\Delta(S) = \bigcup_{n \in S} \Delta(n)$



# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Definition

The delta set of  $S$  is the union  $\Delta(S) = \bigcup_{n \in S} \Delta(n)$

## Goal

Given  $n_1, \dots, n_k$  as input, compute  $\Delta(S)$ .

# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$

$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Definition

The delta set of  $S$  is the union  $\Delta(S) = \bigcup_{n \in S} \Delta(n)$

## Goal

Given  $n_1, \dots, n_k$  as input, compute  $\Delta(S)$ .

The primary difficulty: this is an infinite union!

# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$
$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$
$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Definition

The delta set of  $S$  is the union  $\Delta(S) = \bigcup_{n \in S} \Delta(n)$

## Goal

Given  $n_1, \dots, n_k$  as input, compute  $\Delta(S)$ .

The primary difficulty: this is an infinite union!

## Example

$S = \langle 17, 33, 53, 71 \rangle$ ,  $\Delta(S) = \{2, 4, 6\}$ .

# Delta set of a semigroup

Fix an affine semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$
$$\mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k \quad \rightsquigarrow \quad |\mathbf{a}| = a_1 + \dots + a_k$$
$$L(n) = \{\ell_1 < \dots < \ell_r\} \quad \Delta(n) = \{\ell_i - \ell_{i-1}\}$$

## Definition

The delta set of  $S$  is the union  $\Delta(S) = \bigcup_{n \in S} \Delta(n)$

## Goal

Given  $n_1, \dots, n_k$  as input, compute  $\Delta(S)$ .

The primary difficulty: this is an infinite union!

## Example

$S = \langle 17, 33, 53, 71 \rangle$ ,  $\Delta(S) = \{2, 4, 6\}$ .  $6 \in \Delta(266), \Delta(283), \Delta(300)$ .

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

*For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .*


# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

Example:  $S = \langle 6, 9, 20 \rangle$ :  $2kn_2n_k^2 = 21600$



# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

*For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .*



# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

*For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .*

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$Z(n) = \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\}$$

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \end{aligned}$$

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} \rightarrow Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \leftarrow \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

# The first breakthrough: numerical semigroups

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} \rightarrow Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \leftarrow \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

$$|Z(n)| \approx n^{k-1}$$

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (Chapman–Hoyer–Kaplan, 2009)

For  $n \geq 2kn_2n_k^2$ ,  $\Delta(n) = \Delta(n + n_1n_k)$ .



# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

$S = \langle 6, 9, 20 \rangle$ :  $2kn_2n_k^2 = 21600$ ,  $N_S = 144$ , actual = 91

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

$S = \langle 6, 9, 20 \rangle$ :  $2kn_2n_k^2 = 21600$ ,  $N_S = 144$ , actual = 91

| $S$  | $2kn_2n_k^2$      | $N_S$   | $\Delta(S)$              | Runtime  |
|--|-------------------|---------|--------------------------|----------|
| $\langle 7, 15, 17, 18, 20 \rangle$            | 60000             | 1935    | $\{1, 2, 3\}$            | 1m 28s   |
| $\langle 11, 53, 73, 87 \rangle$               | 3209256           | 14381   | $\{2, 4, 6, 8, 10, 22\}$ | 0m 49s   |
| $\langle 31, 73, 77, 87, 91 \rangle$           | 6045130           | 31364   | $\{2, 4, 6\}$            | 400m 12s |
| $\langle 100, 121, 142, 163, 284 \rangle$      | $9 \cdot 10^7$    | 24850   | ————                     | ————     |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$ | $6 \cdot 10^{15}$ | 2063141 | ————                     | ————     |

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

**Theorem** (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

$S = \langle 6, 9, 20 \rangle$ :  $2kn_2n_k^2 = 21600$ ,  $N_S = 144$ , actual = 91

| $S$  | $2kn_2n_k^2$      | $N_S$   | $\Delta(S)$              | Runtime  |
|--|-------------------|---------|--------------------------|----------|
| $\langle 7, 15, 17, 18, 20 \rangle$            | 60000             | 1935    | $\{1, 2, 3\}$            | 1m 28s   |
| $\langle 11, 53, 73, 87 \rangle$               | 3209256           | 14381   | $\{2, 4, 6, 8, 10, 22\}$ | 0m 49s   |
| $\langle 31, 73, 77, 87, 91 \rangle$           | 6045130           | 31364   | $\{2, 4, 6\}$            | 400m 12s |
| $\langle 100, 121, 142, 163, 284 \rangle$      | $9 \cdot 10^7$    | 24850   | ————                     | ————     |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$ | $6 \cdot 10^{15}$ | 2063141 | ————                     | ————     |

GAP Numerical Semigroups Package, available at

<http://www.gap-system.org/Packages/numericalsgps.html>.

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

# An improvement: the first *usable* algorithm

Fix a **numerical** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}$ .

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq 2kn_2n_k^2 + n_1n_k$ ,  
compute:

$$\begin{aligned} \rightarrow Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1n_1 + \dots + a_kn_k\} \leftarrow \\ Z(n) &\rightsquigarrow L(n) = \{a_1 + \dots + a_k : \mathbf{a} \in Z(n)\} \\ L(n) &= \{\ell_1 < \dots < \ell_r\} \rightsquigarrow \Delta(n) = \{\ell_i - \ell_{i-1}\} \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .



# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ .

## A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$


$S = \langle 6, 9, 20 \rangle$ :

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$S = \langle 6, 9, 20 \rangle$ :




# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$S = \langle 6, 9, 20 \rangle$ :




# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$S = \langle 6, 9, 20 \rangle$ :




# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$S = \langle 6, 9, 20 \rangle$ :






# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$S = \langle 6, 9, 20 \rangle$ :



# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $Z(n)$           | $L(n)$  |
|--------------------------------------|------------------|---------|
| 0                                    | $\{\mathbf{0}\}$ | $\{0\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$        | $Z(n)$             | $L(n)$  |
|---|--------------------|---------|
| 0   | $\{\mathbf{0}\}$   | $\{0\}$ |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$ | $\{\mathbf{e}_1\}$ | $\{1\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$        | $Z(n)$             | $L(n)$  |
|---|--------------------|---------|
| 0   | $\{\mathbf{0}\}$   | $\{0\}$ |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$ | $\{\mathbf{e}_1\}$ | $\{1\}$ |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$ | $\{\mathbf{e}_2\}$ | $\{1\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$            | $Z(n)$              | $L(n)$  |
|---|---------------------|---------|
| 0   | $\{\mathbf{0}\}$    | $\{0\}$ |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$     | $\{\mathbf{e}_1\}$  | $\{1\}$ |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$     | $\{\mathbf{e}_2\}$  | $\{1\}$ |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$ | $\{2\mathbf{e}_1\}$ | $\{2\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ |  | $Z(n)$              | $L(n)$  |
|--------------------------------------|--|---------------------|---------|
| 0                                    |  | $\{\mathbf{0}\}$    | $\{0\}$ |
| 6                                    | $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$    | $\{\mathbf{e}_1\}$  | $\{1\}$ |
| 9                                    | $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$    | $\{\mathbf{e}_2\}$  | $\{1\}$ |
| 12                                   | $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$ | $\{2\mathbf{e}_1\}$ | $\{2\}$ |
| 15                                   | $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$     | $\{(1, 1, 0)\}$     | $\{2\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$            | $Z(n)$              | $L(n)$  |
|---|---------------------|---------|
| 0   | $\{\mathbf{0}\}$    | $\{0\}$ |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$     | $\{\mathbf{e}_1\}$  | $\{1\}$ |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$     | $\{\mathbf{e}_2\}$  | $\{1\}$ |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$ | $\{2\mathbf{e}_1\}$ | $\{2\}$ |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$     | $\{(1, 1, 0)\}$     | $\{2\}$ |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$        |                     |         |



# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$  | $Z(n)$                             | $L(n)$     |
|---|------------------------------------|------------|
| 0   | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$   | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$   | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$   | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$<br>$\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$ | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$  | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned}\phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i\end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$             | $Z(n)$                             | $L(n)$     |
|--|------------------------------------|------------|
| 0  | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$      | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$      | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$      | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$         |                                    |            |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |
| $\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$     |                                    |            |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$             | $Z(n)$                             | $L(n)$     |
|--|------------------------------------|------------|
| 0  | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$      | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$      | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$      | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$         |                                    |            |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |
| $\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$     |                                    |            |
| 20 $\mathbf{0} \xrightarrow{20} \mathbf{e}_3$    | $\{\mathbf{e}_3\}$                 | $\{1\}$    |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$             | $Z(n)$                             | $L(n)$     |
|--|------------------------------------|------------|
| 0  | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$      | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$      | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$      | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$         |                                    |            |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |
| $\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$     |                                    |            |
| 20 $\mathbf{0} \xrightarrow{20} \mathbf{e}_3$    | $\{\mathbf{e}_3\}$                 | $\{1\}$    |
| ⋮  | ⋮                                  | ⋮          |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) & \psi_i : L(n - n_i) &\longrightarrow L(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$             | $Z(n)$                             | $L(n)$           |
|--|------------------------------------|------------------|
| 0  | $\{\mathbf{0}\}$                   | $\{\mathbf{0}\}$ |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$      | $\{\mathbf{e}_1\}$                 | $\{1\}$          |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$      | $\{\mathbf{e}_2\}$                 | $\{1\}$          |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$          |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$      | $\{(1, 1, 0)\}$                    | $\{2\}$          |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$         |                                    |                  |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$       |
| $\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$     |                                    |                  |
| 20 $\mathbf{0} \xrightarrow{20} \mathbf{e}_3$    | $\{\mathbf{e}_3\}$                 | $\{1\}$          |
| ⋮  | ⋮                                  | ⋮                |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) & \qquad \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i & \qquad \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$             | $Z(n)$                             | $L(n)$     |
|--|------------------------------------|------------|
| 0  | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$      | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$      | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$      | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| $\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$         |                                    |            |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |
| $\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$     |                                    |            |
| 20 $\mathbf{0} \xrightarrow{20} \mathbf{e}_3$    | $\{\mathbf{e}_3\}$                 | $\{1\}$    |
| ⋮  | ⋮                                  | ⋮          |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$   | $Z(n)$                             | $L(n)$     |
|--|------------------------------------|------------|
| 0  | $\{\mathbf{0}\}$                   | $\{0\}$    |
| 6 $\mathbf{0} \xrightarrow{6} \mathbf{e}_1$  | $\{\mathbf{e}_1\}$                 | $\{1\}$    |
| 9 $\mathbf{0} \xrightarrow{9} \mathbf{e}_2$  | $\{\mathbf{e}_2\}$                 | $\{1\}$    |
| 12 $\mathbf{e}_1 \xrightarrow{6} 2\mathbf{e}_1$  | $\{2\mathbf{e}_1\}$                | $\{2\}$    |
| 15 $\mathbf{e}_2 \xrightarrow{6} (1, 1, 0)$<br>$\mathbf{e}_1 \xrightarrow{9} (1, 1, 0)$          | $\{(1, 1, 0)\}$                    | $\{2\}$    |
| 18 $2\mathbf{e}_1 \xrightarrow{6} 3\mathbf{e}_1$<br>$\mathbf{e}_2 \xrightarrow{9} 2\mathbf{e}_2$ | $\{3\mathbf{e}_1, 2\mathbf{e}_2\}$ | $\{2, 3\}$ |
| 20 $\mathbf{0} \xrightarrow{20} \mathbf{e}_3$  | $\{\mathbf{e}_3\}$                 | $\{1\}$    |
| $\vdots$   | $\vdots$                           | $\vdots$   |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |
|--------------------------------------|--------|
| 0                                    | {0}    |
| 6                                    |        |
| 9                                    |        |
| 12                                   |        |
| 15                                   |        |
| 18                                   |        |
| 20                                   |        |
| ⋮                                    |        |



# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$                                  |
|--------------------------------------|---|
| 0                                    | {0}                                     |
| 6                                    | {1} $0 \overset{6}{\rightsquigarrow} 1$ |
| 9                                    |   |
| 12                                   |   |
| 15                                   |   |
| 18                                   |   |
| 20                                   |   |
| ⋮                                    |   |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                       |
|--------------------------------------|--------|-----------------------|
| 0                                    | {0}    |                       |
| 6                                    | {1}    | $0 \xrightarrow{6} 1$ |
| 9                                    | {1}    | $0 \xrightarrow{9} 1$ |
| 12                                   |        |                       |
| 15                                   |        |                       |
| 18                                   |        |                       |
| 20                                   |        |                       |
| ⋮                                    |        |                       |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i \end{aligned}$$

$$\begin{aligned} \psi_i : L(n - n_i) &\longrightarrow L(n) \\ \ell &\longmapsto \ell + 1 \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                       |
|--------------------------------------|--------|-----------------------|
| 0                                    | {0}    |                       |
| 6                                    | {1}    | $0 \xrightarrow{6} 1$ |
| 9                                    | {1}    | $0 \xrightarrow{9} 1$ |
| 12                                   | {2}    | $1 \xrightarrow{6} 2$ |
| 15                                   |        |                       |
| 18                                   |        |                       |
| 20                                   |        |                       |
| ⋮                                    |        |                       |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                                     |
|--------------------------------------|--------|-------------------------------------|
| 0                                    | {0}    |                                     |
| 6                                    | {1}    | $0 \overset{6}{\rightsquigarrow} 1$ |
| 9                                    | {1}    | $0 \overset{9}{\rightsquigarrow} 1$ |
| 12                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$ |
| 15                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$ |
| 18                                   |        |                                     |
| 20                                   |        |                                     |
| ⋮                                    |        |                                     |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) & \psi_i : L(n - n_i) &\longrightarrow L(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i & \ell &\longmapsto \ell + 1 \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                       |
|--------------------------------------|--------|-----------------------|
| 0                                    | {0}    |                       |
| 6                                    | {1}    | $0 \xrightarrow{6} 1$ |
| 9                                    | {1}    | $0 \xrightarrow{9} 1$ |
| 12                                   | {2}    | $1 \xrightarrow{6} 2$ |
| 15                                   | {2}    | $1 \xrightarrow{6} 2$ |
|                                      |        | $1 \xrightarrow{9} 2$ |
| 18                                   |        |                       |
| 20                                   |        |                       |
| ⋮                                    |        |                       |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                       |
|--------------------------------------|--------|-----------------------|
| 0                                    | {0}    |                       |
| 6                                    | {1}    | $0 \xrightarrow{6} 1$ |
| 9                                    | {1}    | $0 \xrightarrow{9} 1$ |
| 12                                   | {2}    | $1 \xrightarrow{6} 2$ |
| 15                                   | {2}    | $1 \xrightarrow{6} 2$ |
|                                      |        | $1 \xrightarrow{9} 2$ |
| 18                                   | {2, 3} | $2 \xrightarrow{6} 3$ |
| 20                                   |        |                       |
| ⋮                                    |        |                       |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{aligned} \phi_i : Z(n - n_i) &\longrightarrow Z(n) & \psi_i : L(n - n_i) &\longrightarrow L(n) \\ \mathbf{a} &\longmapsto \mathbf{a} + \mathbf{e}_i & \ell &\longmapsto \ell + 1 \end{aligned}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                                     |
|--------------------------------------|--------|-------------------------------------|
| 0                                    | {0}    |                                     |
| 6                                    | {1}    | $0 \overset{6}{\rightsquigarrow} 1$ |
| 9                                    | {1}    | $0 \overset{9}{\rightsquigarrow} 1$ |
| 12                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$ |
| 15                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$ |
|                                      |        | $1 \overset{9}{\rightsquigarrow} 2$ |
| 18                                   | {2, 3} | $2 \overset{6}{\rightsquigarrow} 3$ |
|                                      |        | $1 \overset{9}{\rightsquigarrow} 2$ |
| 20                                   |        |                                     |
| ⋮                                    |        |                                     |

# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$ |                                      |
|--------------------------------------|--------|--------------------------------------|
| 0                                    | {0}    |                                      |
| 6                                    | {1}    | $0 \overset{6}{\rightsquigarrow} 1$  |
| 9                                    | {1}    | $0 \overset{9}{\rightsquigarrow} 1$  |
| 12                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$  |
| 15                                   | {2}    | $1 \overset{6}{\rightsquigarrow} 2$  |
|                                      |        | $1 \overset{9}{\rightsquigarrow} 2$  |
| 18                                   | {2, 3} | $2 \overset{6}{\rightsquigarrow} 3$  |
|                                      |        | $1 \overset{9}{\rightsquigarrow} 2$  |
| 20                                   | {1}    | $0 \overset{20}{\rightsquigarrow} 1$ |
| ⋮                                    |        |                                      |



# A faster solution: dynamic programming

Fix  $n \in S = \langle n_1, \dots, n_k \rangle$ . For each  $i \leq k$ ,

$$\begin{array}{ccc} \phi_i : Z(n - n_i) & \longrightarrow & Z(n) \\ \mathbf{a} & \longmapsto & \mathbf{a} + \mathbf{e}_i \end{array} \qquad \begin{array}{ccc} \psi_i : L(n - n_i) & \longrightarrow & L(n) \\ \ell & \longmapsto & \ell + 1 \end{array}$$

$$Z(n) = \bigcup_{i \leq k} \phi_i(Z(n - n_i))$$

$$L(n) = \bigcup_{i \leq k} \psi_i(L(n - n_i))$$

| $n \in S = \langle 6, 9, 20 \rangle$ | $L(n)$   |                                      |
|--------------------------------------|----------|--------------------------------------|
| 0                                    | {0}      |                                      |
| 6                                    | {1}      | $0 \overset{6}{\rightsquigarrow} 1$  |
| 9                                    | {1}      | $0 \overset{9}{\rightsquigarrow} 1$  |
| 12                                   | {2}      | $1 \overset{6}{\rightsquigarrow} 2$  |
| 15                                   | {2}      | $1 \overset{6}{\rightsquigarrow} 2$  |
|                                      |          | $1 \overset{9}{\rightsquigarrow} 2$  |
| 18                                   | {2, 3}   | $2 \overset{6}{\rightsquigarrow} 3$  |
|                                      |          | $1 \overset{9}{\rightsquigarrow} 2$  |
| 20                                   | {1}      | $0 \overset{20}{\rightsquigarrow} 1$ |
| $\vdots$                             | $\vdots$ | $\vdots$                             |

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq N_S + \text{lcm}(n_1, n_k)$ ,  
compute:

$$\begin{aligned} Z(n) &= \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k\} \\ Z(n) &\rightsquigarrow L(n) \\ L(n) &\rightsquigarrow \Delta(n) \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq N_S + \text{lcm}(n_1, n_k)$ ,  
compute:

$$\begin{aligned} Z(n) &= \{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \} \\ L(n - *) &\rightsquigarrow L(n) \\ L(n) &\rightsquigarrow \Delta(n) \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq N_S + \text{lcm}(n_1, n_k)$ ,  
compute:

$$\begin{aligned} \cancel{Z(n)} &= \{ \cancel{\mathbf{a}} \in \mathbb{Z}_{\geq 0}^k : n = \cancel{a_1 n_1} + \dots + \cancel{a_k n_k} \} \\ &\quad L(n - *) \rightsquigarrow L(n) \\ &\quad L(n) \rightsquigarrow \Delta(n) \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

This is *significantly* faster!

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq N_S + \text{lcm}(n_1, n_k)$ ,  
compute:

$$\begin{aligned} Z(n) &= \{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \} \\ L(n - *) &\rightsquigarrow L(n) \\ L(n) &\rightsquigarrow \Delta(n) \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

This is *significantly* faster!

$$|Z(n)| \approx n^{k-1}$$

# Computing the delta set dynamically

Theorem (García-García–Moreno-Frías–Vigneron-Tenorio, 2014)

$S = \langle n_1, \dots, n_k \rangle$ . For  $n \geq N_S$ ,  $\Delta(n) = \Delta(n + \text{lcm}(n_1, n_k))$ .

For  $n \in S$  with  $0 \leq n \leq N_S + \text{lcm}(n_1, n_k)$ ,  
compute:

$$\begin{aligned} Z(n) &= \{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \} \\ L(n - *) &\rightsquigarrow L(n) \\ L(n) &\rightsquigarrow \Delta(n) \end{aligned}$$

Compute  $\Delta(S) = \bigcup_n \Delta(n)$ .

This is *significantly* faster!

$$\begin{aligned} |Z(n)| &\approx n^{k-1} \\ |L(n)| &\approx n \end{aligned}$$

# Runtime comparison



# Runtime comparison

| $S$  | $N_S$   | $\Delta(S)$              | Manual   | Dynamic |
|--|---------|--------------------------|----------|---------|
| $\langle 7, 15, 17, 18, 20 \rangle$            | 1935    | $\{1, 2, 3\}$            | 1m 28s   | 146ms   |
| $\langle 11, 53, 73, 87 \rangle$               | 14381   | $\{2, 4, 6, 8, 10, 22\}$ | 0m 49s   | 2.5s    |
| $\langle 31, 73, 77, 87, 91 \rangle$           | 31364   | $\{2, 4, 6\}$            | 400m 12s | 4.2s    |
| $\langle 100, 121, 142, 163, 284 \rangle$      | 24850   | $\{21\}$                 | ————     | 0m 3.6s |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$ | 2063141 | $\{10, 20, 30\}$         | ————     | 1m 56s  |

# Runtime comparison

| $S$  | $N_S$   | $\Delta(S)$              | Manual   | Dynamic |
|--|---------|--------------------------|----------|---------|
| $\langle 7, 15, 17, 18, 20 \rangle$            | 1935    | $\{1, 2, 3\}$            | 1m 28s   | 146ms   |
| $\langle 11, 53, 73, 87 \rangle$               | 14381   | $\{2, 4, 6, 8, 10, 22\}$ | 0m 49s   | 2.5s    |
| $\langle 31, 73, 77, 87, 91 \rangle$           | 31364   | $\{2, 4, 6\}$            | 400m 12s | 4.2s    |
| $\langle 100, 121, 142, 163, 284 \rangle$      | 24850   | $\{21\}$                 | ————     | 0m 3.6s |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$ | 2063141 | $\{10, 20, 30\}$         | ————     | 1m 56s  |

GAP Numerical Semigroups Package, available at

<http://www.gap-system.org/Packages/numericalsgps.html>.

# Generalize to affine semigroups?

Key obstruction: what does “eventually periodic” mean?

# Generalize to affine semigroups?



Key obstruction: what does “eventually periodic” mean?

Example:  $S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$

# Generalize to affine semigroups?

Key obstruction: what does “eventually periodic” mean?

Example:  $S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$





$$\Delta(S) = \{1, 2, 4\}$$

# Generalize to affine semigroups?

Key obstruction: what does “eventually periodic” mean?

Example:  $S = \langle (1, 1), (1, 5), (2, 5), (3, 5), (5, 1), (5, 2), (5, 3) \rangle \subset \mathbb{Z}_{\geq 0}^2$



$$\Delta(S) = \{1, 2, 4\}$$

Need a new approach!

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \longleftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \longleftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\begin{aligned} \mathbf{a} &\sim \mathbf{a} \\ \mathbf{a} \sim \mathbf{b} &\Rightarrow \mathbf{b} \sim \mathbf{a} \\ \mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} &\Rightarrow \mathbf{a} \sim \mathbf{c} \end{aligned}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*.

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

Monomial map:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a} \quad x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$\begin{aligned} x^{\mathbf{a}} - x^{\mathbf{a}} &= 0 \in I_S \\ x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S &\Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S \end{aligned}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S$$

$$(x^{\mathbf{a}} - x^{\mathbf{b}}) + (x^{\mathbf{b}} - x^{\mathbf{c}}) = x^{\mathbf{a}} - x^{\mathbf{c}}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Definition

The *kernel*  $\ker \pi$  is the relation  $\sim$  on  $\mathbb{Z}_{\geq 0}^k$  with  $\mathbf{a} \sim \mathbf{b}$  whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$  is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S$$

$$(x^{\mathbf{a}} - x^{\mathbf{b}}) + (x^{\mathbf{b}} - x^{\mathbf{c}}) = x^{\mathbf{a}} - x^{\mathbf{c}}$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{c}}(x^{\mathbf{a}} - x^{\mathbf{b}}) \in I_S$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle:$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

$Z(18)$ :

$(3, 0, 0)$



$(0, 2, 0)$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

$$Z(18): \quad Z(60): \quad \begin{array}{c} \bullet (10, 0, 0) \\ \bullet (7, 2, 0) \\ \bullet (4, 4, 0) \\ \bullet (1, 6, 0) \end{array} \quad x^7 y^2 - x^4 y^4 = x^4 y^2 (x^3 - y^2)$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:



$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

|   |           |  |   |
|---|-----------|--|---|
| Z(18):  | Z(60):    |  | $x^7 y^2 - x^4 y^4 = x^4 y^2 (x^3 - y^2)$ |
| (3, 0, 0)   |           |  | $x^7 y^2 - z^3 = (x^7 y^2 - x^4 y^4)$     |
|  | (0, 0, 3) | (4, 4, 0)  | $+ (x^4 y^4 - z^3)$                       |
| (0, 2, 0)   |           | (1, 6, 0)  |   |

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



$$\begin{aligned} x^7 y^2 - x^4 y^4 &= x^4 y^2 (x^3 - y^2) \\ x^7 y^2 - z^3 &= (x^7 y^2 - x^4 y^4) \\ &\quad + (x^4 y^4 - z^3) \end{aligned}$$

Generating set for  $I_S \iff \pi^{-1}(n)$  connected for all  $n \in S$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

$Z(18)$ :

$$(3, 0, 0)$$



$$(0, 2, 0)$$

$Z(60)$ :

$$(10, 0, 0)$$



$$(7, 2, 0)$$

$$(4, 4, 0)$$

$$(1, 6, 0)$$

$$(0, 0, 3)$$

All minimal generating sets:



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



All minimal generating sets:

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



All minimal generating sets:

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



All minimal generating sets:

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



All minimal generating sets:

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$



# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$

$\mathbb{Z}(18)$ :

$$(3, 0, 0)$$



$$(0, 2, 0)$$

$\mathbb{Z}(60)$ :

$$\bullet (10, 0, 0)$$

$$\bullet (7, 2, 0)$$

$$\bullet (4, 4, 0)$$

$$\bullet (1, 6, 0)$$

$$(0, 0, 3)$$



All minimal generating sets:

$$I_S = \langle x^3 - y^2, x^{10} - z^3 \rangle$$

$$= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle$$

$$= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle$$

$$= \langle x^3 - y^2, x^6 y - z^3 \rangle$$

# Commutative algebra hiding in the background

Fix an **affine** semigroup  $S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d$ .

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:


$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

## Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$$



All minimal generating sets:

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$

# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$



$Z(550)$

●  $(2, 1, 0, 4)$

●  $(22, 6, 0, 0)$

●  $(6, 8, 0, 1)$

●  $(24, 3, 1, 0)$

●  $(8, 5, 1, 1)$

●  $(26, 0, 2, 0)$



●  $(10, 2, 2, 1)$

# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \quad (\text{affine}) \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$







# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \quad (\text{affine}) \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$





# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$





# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$





# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \quad (\text{affine}) \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$





# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \quad (\text{affine}) \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$






# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \quad (\text{affine}) \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$






# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$






# Commutative algebra hiding in the background

$$S = \langle n_1, \dots, n_k \rangle \subset \mathbb{Z}_{\geq 0}^d \text{ (affine)} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example:  $S = \langle 13, 44, 106, 120 \rangle$ .

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$



# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$
$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

Idea: only connect *some* of the factorizations

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a} - \mathbf{b}\| \leq j \rangle \subset I_S$$

Idea: only connect *some* of the factorizations

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a} - \mathbf{b}\| \leq j \rangle \subset I_S$$

Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

# The delta set via commutative algebra

$$\begin{array}{ccc} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k \end{array} \qquad \begin{array}{ccc} \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k \end{array} \qquad \begin{array}{lll} \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 28



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 21



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 21



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 2



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 2



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 2



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 2



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: **1**





# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 1



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S \langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 1



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 1



# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$


Idea: only connect *some* of the factorizations

$$I_0 \subset I_1 \subset I_2 \subset I_3 \subset I_4 \subset \dots \subset I_S$$

Example:  $S \langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

$Z(244)$ :

connected components: 1



$$I_0 = \langle x^{11} z^3 - y^{14} \rangle$$

$$I_1 = I_0 + \langle x^3 - y^2, x^8 z^3 - y^{12} \rangle$$

$$I_2 = I_1 + \langle x^5 z^3 - y^{10} \rangle$$

$$I_3 = I_2 + \langle x^2 z^3 - y^8 \rangle$$

$$I_4 = I_3 + \langle x^4 y^4 - z^3 \rangle$$

$$= I_5 = I_6 = \dots = I_S$$

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

## Theorem (O, 2016)

*In the ascending chain  $I_0 \subset I_1 \subset I_2 \subset \dots \subset I_S$ ,*

$$j \in \Delta(S) \quad \text{if and only if} \quad I_{j-1} \subsetneq I_j$$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

## Theorem (O, 2016)

In the ascending chain  $I_0 \subset I_1 \subset I_2 \subset \dots \subset I_S$ ,

$$j \in \Delta(S) \quad \text{if and only if} \quad I_{j-1} \subsetneq I_j$$

Algorithm for computing  $\Delta(S)$ :

- Compute generators for  $I_0, I_1, \dots$
- At each step, check if  $I_{j-1} \neq I_j$
- Stop when  $I_S$  reached

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$



# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

$I_{\text{hom}}$ : homogenization of  $I_S$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \quad \longrightarrow \quad x^{\mathbf{a}} - t^{|\mathbf{a}|-|\mathbf{b}|} x^{\mathbf{b}} \in I_{\text{hom}}$$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

$I_{\text{hom}}$ : homogenization of  $I_S$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \quad \longrightarrow \quad x^{\mathbf{a}} - t^{|\mathbf{a}|-|\mathbf{b}|} x^{\mathbf{b}} \in I_{\text{hom}}$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, xy^6 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

$I_{\text{hom}}$ : homogenization of  $I_S$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \quad \longrightarrow \quad x^{\mathbf{a}} - t^{|\mathbf{a}|-|\mathbf{b}|} x^{\mathbf{b}} \in I_{\text{hom}}$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, xy^6 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

Lex Gröbner basis for  $I_{\text{hom}}$ :

$$\begin{aligned} I_{\text{hom}} = \langle & x^{11}z^3 - y^{14}, \\ & x^3 - ty^2, x^8z^3 - ty^{12}, \\ & t^2x^5z^3 - y^{10}, \\ & t^3x^2z^3 - y^8, \\ & xy^6 - t^4z^3 \rangle \end{aligned}$$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

$I_{\text{hom}}$ : homogenization of  $I_S$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \quad \longrightarrow \quad x^{\mathbf{a}} - t^{|\mathbf{a}|-|\mathbf{b}|} x^{\mathbf{b}} \in I_{\text{hom}}$$

Example:  $S\langle 6, 9, 20 \rangle \quad I_S = \langle x^3 - y^2, xy^6 - z^3 \rangle \subset \mathbb{k}[x, y, z]$

Lex Gröbner basis for  $I_{\text{hom}}$ :

$$\begin{array}{ll} I_{\text{hom}} = \langle x^{11}z^3 - y^{14}, & I_0 = \langle x^{11}z^3 - y^{14} \rangle \\ x^3 - ty^2, x^8z^3 - ty^{12}, & I_1 = I_0 + \langle x^3 - y^2, x^8z^3 - y^{12} \rangle \\ t^2x^5z^3 - y^{10}, & I_2 = I_1 + \langle x^5z^3 - y^{10} \rangle \\ t^3x^2z^3 - y^8, & I_3 = I_2 + \langle x^2z^3 - y^8 \rangle \\ xy^6 - t^4z^3 \rangle & I_4 = I_3 + \langle xy^6 - z^3 \rangle \end{array}$$

# The delta set via commutative algebra

$$\begin{array}{llll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k & x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

# The delta set via commutative algebra

$$\begin{array}{lll} \pi : \mathbb{Z}_{\geq 0}^k & \longrightarrow & S = \langle n_1, \dots, n_k \rangle \\ \mathbf{a} & \longmapsto & a_1 n_1 + \dots + a_k n_k \end{array} \qquad \begin{array}{lll} \varphi : \mathbb{k}[x_1, \dots, x_k] & \longrightarrow & \mathbb{k}[w] \\ x_i & \longmapsto & w^{n_i} \end{array}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

Algorithm to compute  $\Delta(S)$  (García-Sánchez-O-Webb, 2018)

- Homogenize the ideal  $I_S$  with a new variable  $t$
- Compute a reduced lex Gröbner basis  $G$  with  $t < x_i$
- $\Delta(S) = \{d : t^d x^{\mathbf{a}} - x^{\mathbf{b}} \in G\}$

# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

## Algorithm to compute $\Delta(S)$ (García-Sánchez-O-Webb, 2018)

- Homogenize the ideal  $I_S$  with a new variable  $t$
- Compute a reduced lex Gröbner basis  $G$  with  $t < x_i$
- $\Delta(S) = \{d : t^d x^{\mathbf{a}} - x^{\mathbf{b}} \in G\}$

| $S$  | $\Delta(S)$      | Manual | Dynamic | Algebraic |
|--|------------------|--------|---------|-----------|
| $\langle 100, 121, 142, 163, 284 \rangle$      | $\{21\}$         | Days   | 0m 3.6s | $< 10$ ms |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$ | $\{10, 20, 30\}$ | Days   | 1m 56s  | $< 10$ ms |

# The delta set via commutative algebra

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow S = \langle n_1, \dots, n_k \rangle & \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k & x_i &\longmapsto w^{n_i} \end{aligned}$$

$$I_S = \ker(\varphi) = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \rangle$$

$$I_j = \langle x^{\mathbf{a}} - x^{\mathbf{b}} : \pi(\mathbf{a}) = \pi(\mathbf{b}) \text{ and } \|\mathbf{a}\| - \|\mathbf{b}\| \leq j \rangle \subset I_S$$

## Algorithm to compute $\Delta(S)$ (García-Sánchez-O-Webb, 2018)

- Homogenize the ideal  $I_S$  with a new variable  $t$
- Compute a reduced lex Gröbner basis  $G$  with  $t < x_i$
- $\Delta(S) = \{d : t^d x^{\mathbf{a}} - x^{\mathbf{b}} \in G\}$

| $S$   | $\Delta(S)$   | Manual | Dynamic | Algebraic |
|---|---|--------|---------|-----------|
| $\langle 100, 121, 142, 163, 284 \rangle$   | $\{21\}$  | Days   | 0m 3.6s | < 10 ms   |
| $\langle 1001, 1211, 1421, 1631, 2841 \rangle$  | $\{10, 20, 30\}$  | Days   | 1m 56s  | < 10 ms   |
| $\langle 550, 1060, 1600, 1781, 4126, 4139, 4407, 5167, 6073, 6079, 6169, 7097, 7602, 8782, 8872 \rangle$ | $\left\{ \begin{array}{l} 1, 2, 3, 4, 5, 6, 7, \\ 8, 9, 10, 11, 12, 13, \\ 14, 15, 16, 17, 19 \end{array} \right\}$ | Years  | Days    | < 1 min   |



# References



T. Barron, C. O'Neill, R. Pelayo (2015)

On the computation of delta sets and  $\omega$ -primality in numerical monoids.  
preprint.



J. García-García, M. Moreno-Frías, A. Vigneron-Tenorio (2014)

Computation of delta sets of numerical monoids.  
preprint.



P. García-Sánchez, C. O'Neill, G. Webb (2017)

On the computation of factorization invariants for affine semigroups.  
preprint.



M. Delgado, P. García-Sánchez, J. Morais

GAP Numerical Semigroups Package

<http://www.gap-system.org/Packages/numericalsgps.html>.

# References



T. Barron, C. O'Neill, R. Pelayo (2015)

On the computation of delta sets and  $\omega$ -primality in numerical monoids.  
preprint.



J. García-García, M. Moreno-Frías, A. Vigneron-Tenorio (2014)

Computation of delta sets of numerical monoids.  
preprint.



P. García-Sánchez, C. O'Neill, G. Webb (2017)

On the computation of factorization invariants for affine semigroups.  
preprint.



M. Delgado, P. García-Sánchez, J. Morais

GAP Numerical Semigroups Package

<http://www.gap-system.org/Packages/numericalsgps.html>.

Thanks!