

Numerical semigroups, minimal presentations, and posets

Christopher O'Neill

San Diego State University

cdoneill@sdsu.edu

Joint with (i) Winfred Bruns, Pedro García-Sánchez, and Dane Wilbourne;
(ii) Nathan Kaplan; (iii) *Tara Gomes and *Eduardo Torres Davila;
(iv) *Ceyhun Elmacioglu, Kieran Hilmer, *Hannah Koufmann, and *Melin Okandan

* = undergraduate student

Slides available: <https://cdoneill.sdsu.edu/>

January 23, 2025

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example:

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Example: $S = \langle 6, 9, 18, 20, 32 \rangle$

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Example: $S = \langle 6, 9, \del{18}, 20, \del{32} \rangle$

Definition

A *numerical semigroup* $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Example: $S = \langle 6, 9, \del{18}, 20, \del{32} \rangle = McN$

Numerical semigroups

Definition

A numerical semigroup $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Example: $S = \langle 6, 9, \del{18}, 20, \del{32} \rangle = McN$

Fact

Every numerical semigroup has a unique minimal generating set.

Numerical semigroups

Definition

A numerical semigroup $S \subseteq \mathbb{Z}_{\geq 0}$: closed under **addition**, $|\mathbb{Z}_{\geq 0} \setminus S| < \infty$.

Example: “McNugget Semigroup”

$$McN = \langle 6, 9, 20 \rangle = \left\{ \begin{array}{l} 0, 6, 9, 12, 15, 18, 20, 21, 24, \dots \\ \dots, 36, 38, 39, 40, 41, 42, 44 \rightarrow \end{array} \right\}$$

Example: $S = \langle 6, 9, \del{18}, 20, \del{32} \rangle = McN$

Fact

Every numerical semigroup has a unique minimal generating set.

Multiplicity: $m(S) =$ smallest nonzero element

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Observations:

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Observations:

- The elements of $\text{Ap}(S)$ are distinct modulo m

Apéry sets

Fix a numerical semigroup S with $m(S) = m$.

Definition

The *Apéry set* of S is

$$\text{Ap}(S) = \{a \in S : a - m \notin S\}$$

If $S = \langle 6, 9, 20 \rangle$, then

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

For 2 mod 6: $\{2, 8, 14, 20, 26, 32, \dots\} \cap S = \{20, 26, 32, \dots\}$

For 3 mod 6: $\{3, 9, 15, 21, \dots\} \cap S = \{9, 15, 21, \dots\}$

For 4 mod 6: $\{4, 10, 16, 22, \dots\} \cap S = \{40, 46, 52, \dots\}$

Observations:

- The elements of $\text{Ap}(S)$ are distinct modulo m
- $|\text{Ap}(S)| = m$

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

but $a_1 + a_2 \equiv 3 \pmod{5}$ and $a_1 + a_2 < a_3$.

Apéry sets

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

but $a_1 + a_2 \equiv 3 \pmod{5}$ and $a_1 + a_2 < a_3$.

Is $\{0, 13, 14, 27, 10, 11\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 6, \quad a_1 = 13, \quad a_2 = 14, \quad a_3 = 27, \quad a_4 = 10, \quad a_5 = 11$$

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

but $a_1 + a_2 \equiv 3 \pmod{5}$ and $a_1 + a_2 < a_3$.

Is $\{0, 13, 14, 27, 10, 11\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 6, \quad a_1 = 13, \quad a_2 = 14, \quad a_3 = 27, \quad a_4 = 10, \quad a_5 = 11$$

but $a_4 + a_5 \equiv 3 \pmod{6}$ and $a_4 + a_5 < a_3$.

Apéry sets

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

but $a_1 + a_2 \equiv 3 \pmod{5}$ and $a_1 + a_2 < a_3$.

Is $\{0, 13, 14, 27, 10, 11\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 6, \quad a_1 = 13, \quad a_2 = 14, \quad a_3 = 27, \quad a_4 = 10, \quad a_5 = 11$$

but $a_4 + a_5 \equiv 3 \pmod{6}$ and $a_4 + a_5 < a_3$.

Theorem

If $A = \{0, a_1, \dots, a_{m-1}\}$ with each $a_i > m$ and $a_i \equiv i \pmod{m}$, then there exists a numerical semigroup S with $\text{Ap}(S) = A$ if and only if

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

Is $A = \{0, 11, 7, 23, 19\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 5, \quad a_1 = 11, \quad a_2 = 7, \quad a_3 = 23, \quad a_4 = 19$$

but $a_1 + a_2 \equiv 3 \pmod{5}$ and $a_1 + a_2 < a_3$.

Is $\{0, 13, 14, 27, 10, 11\}$ the Apéry set of some numerical semigroup?

$$m = |A| = 6, \quad a_1 = 13, \quad a_2 = 14, \quad a_3 = 27, \quad a_4 = 10, \quad a_5 = 11$$

but $a_4 + a_5 \equiv 3 \pmod{6}$ and $a_4 + a_5 < a_3$.

Theorem

If $A = \{0, a_1, \dots, a_{m-1}\}$ with each $a_i > m$ and $a_i \equiv i \pmod{m}$, then there exists a numerical semigroup S with $\text{Ap}(S) = A$ if and only if

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

Big idea: the inequalities “ $a_i + a_j \geq a_{i+j}$ ” to define a **cone** C_m .

Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Kunz cone


Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Example: C_3




Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Example: C_3



$$S = \langle 3, 5, 7 \rangle$$

$$\text{Ap}(S) = \{0, 7, 5\}$$


Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Example: C_3



$$\begin{aligned} S &= \langle 3, 5, 7 \rangle \\ \text{Ap}(S) &= \{0, 7, 5\} \end{aligned}$$

$$\begin{aligned} S &= \langle 3, 4 \rangle \\ \text{Ap}(S) &= \{0, 4, 8\} \end{aligned}$$

Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Kunz cone


Definition

The *Kunz cone* $C_m \subseteq \mathbb{R}^{m-1}$ is a pointed cone with defining inequalities

$$a_i + a_j \geq a_{i+j} \quad \text{whenever} \quad i + j \neq 0.$$

$$\begin{aligned} \{S \subseteq \mathbb{Z}_{\geq 0} : m(S) = m\} &\longrightarrow C_m \\ \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\} &\longmapsto (a_1, \dots, a_{m-1}) \end{aligned}$$

Example: C_4



Question

When are numerical semigroups in (the relative interior of) the same face?

Question

When are numerical semigroups in (the relative interior of) the same face?

Big picture: “moduli space” approach for studying XYZ 's

- Define a space with XYZ 's as points
Small changes to an $XYZ \rightsquigarrow$ small movements in space
- Let geometric/topological structure inform study of XYZ 's

Question

When are numerical semigroups in (the relative interior of) the same face?

Big picture: “moduli space” approach for studying XYZ 's

- Define a space with XYZ 's as points
Small changes to an $XYZ \rightsquigarrow$ small movements in space
- Let geometric/topological structure inform study of XYZ 's

Basic example: $GL_n(\mathbb{R}) \hookrightarrow \mathbb{R}^{n^2}$

Faces of the Kunz cone

Question


When are numerical semigroups in (the relative interior of) the same face?

Big picture: “moduli space” approach for studying XYZ 's

- Define a space with XYZ 's as points
Small changes to an $XYZ \rightsquigarrow$ small movements in space
- Let geometric/topological structure inform study of XYZ 's

Basic example: $GL_n(\mathbb{R}) \hookrightarrow \mathbb{R}^{n^2}$

More interesting example: C_m



Question

When are numerical semigroups in (the relative interior of) the same face?

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

If $S = \langle n_1, \dots, n_k \rangle$, then

$$n_i \not\equiv n_j \pmod{n_1} \implies k \leq m(S)$$

Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

If $S = \langle n_1, \dots, n_k \rangle$, then

$$n_i \not\equiv n_j \pmod{n_1} \implies k \leq m(S)$$

If $k = m(S)$, then S has *max embedding dimension*

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

If $S = \langle n_1, \dots, n_k \rangle$, then

$$n_i \not\equiv n_j \pmod{n_1} \implies k \leq m(S)$$

If $k = m(S)$, then S has *max embedding dimension*

$$S = \langle m, a_1, \dots, a_{m-1} \rangle \quad \text{where} \quad \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\}$$

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

If $S = \langle n_1, \dots, n_k \rangle$, then

$$n_i \not\equiv n_j \pmod{n_1} \implies k \leq m(S)$$

If $k = m(S)$, then S has *max embedding dimension*

$$S = \langle m, a_1, \dots, a_{m-1} \rangle \quad \text{where} \quad \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\}$$

Geometrically: “most” numerical semigroups with $m(S) = m$ are MED

Question

When are numerical semigroups in (the relative interior of) the same face?

First steps: $S \in \text{Int}(C_m)$ if and only if S has *max embedding dimension*

If $S = \langle n_1, \dots, n_k \rangle$, then

$$n_i \not\equiv n_j \pmod{n_1} \implies k \leq m(S)$$

If $k = m(S)$, then S has *max embedding dimension*

$$S = \langle m, a_1, \dots, a_{m-1} \rangle \quad \text{where} \quad \text{Ap}(S) = \{0, a_1, \dots, a_{m-1}\}$$

Geometrically: “most” numerical semigroups with $m(S) = m$ are MED

What about the other faces?

Question

When are numerical semigroups in (the relative interior of) the same face?

Question

When are numerical semigroups in (the relative interior of) the same face?

Example: $S = \langle 4, 10, 11, 13 \rangle$

$$\text{Ap}(S) = \{0, 13, 10, 11\}$$

$$a_1 = 13, \quad a_2 = 10, \quad a_3 = 11$$

$$2a_1 > a_2 \quad a_1 + a_2 > a_3$$

$$2a_3 > a_2 \quad a_2 + a_3 > a_1$$

Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

Example: $S = \langle 4, 10, 11, 13 \rangle$

$$\text{Ap}(S) = \{0, 13, 10, 11\}$$
$$a_1 = 13, \quad a_2 = 10, \quad a_3 = 11$$

$$2a_1 > a_2 \quad a_1 + a_2 > a_3$$
$$2a_3 > a_2 \quad a_2 + a_3 > a_1$$

Example: $S = \langle 4, 10, 13 \rangle$

$$\text{Ap}(S) = \{0, 13, 10, 23\}$$
$$a_1 = 13, \quad a_2 = 10, \quad a_3 = 23$$

$$2a_1 > a_2 \quad a_1 + a_2 = a_3$$
$$2a_3 > a_2 \quad a_2 + a_3 > a_1$$

Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

Example: $S = \langle 4, 10, 11, 13 \rangle$

$$\begin{array}{l} \text{Ap}(S) = \{0, 13, 10, 11\} \\ a_1 = 13, \quad a_2 = 10, \quad a_3 = 11 \end{array} \qquad \begin{array}{ll} 2a_1 > a_2 & a_1 + a_2 > a_3 \\ 2a_3 > a_2 & a_2 + a_3 > a_1 \end{array}$$

Example: $S = \langle 4, 10, 13 \rangle$

$$\begin{array}{l} \text{Ap}(S) = \{0, 13, 10, 23\} \\ a_1 = 13, \quad a_2 = 10, \quad a_3 = 23 \end{array} \qquad \begin{array}{ll} 2a_1 > a_2 & a_1 + a_2 = a_3 \\ 2a_3 > a_2 & a_2 + a_3 > a_1 \end{array}$$

Example: $S = \langle 4, 13 \rangle$

$$\begin{array}{l} \text{Ap}(S) = \{0, 13, 26, 39\} \\ a_1 = 13, \quad a_2 = 26, \quad a_3 = 39 \end{array} \qquad \begin{array}{ll} 2a_1 = a_2 & a_1 + a_2 = a_3 \\ 2a_3 > a_2 & a_2 + a_3 > a_1 \end{array}$$

Question

When are numerical semigroups in (the relative interior of) the same face?

Faces of the Kunz cone


Question

When are numerical semigroups in (the relative interior of) the same face?

Definition

The *Apéry poset* of S : define $a \preceq a'$ whenever $a' - a \in S$.

$$\text{Ap}(S) = \{0, 13, 10, 23\}$$



$$\text{Ap}(S) = \{0, 13, 26, 39\}$$



Question

When are numerical semigroups in (the relative interior of) the same face?

Question

When are numerical semigroups in (the relative interior of) the same face?

$$S = \langle 6, 9, 20 \rangle$$
$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$

$$S' = \langle 6, 26, 27 \rangle$$
$$\text{Ap}(S') = \{0, 79, 26, 27, 52, 53\}$$


Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?


$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



$$S' = \langle 6, 26, 27 \rangle$$

$$\text{Ap}(S') = \{0, 79, 26, 27, 52, 53\}$$




Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?


$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



$$S' = \langle 6, 26, 27 \rangle$$

$$\text{Ap}(S') = \{0, 79, 26, 27, 52, 53\}$$




Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?


$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



$$S' = \langle 6, 26, 27 \rangle$$

$$\text{Ap}(S') = \{0, 79, 26, 27, 52, 53\}$$



The *Kunz poset* of S : use ground set \mathbb{Z}_m instead of $\text{Ap}(S)$.


Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?


$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



$$S' = \langle 6, 26, 27 \rangle$$

$$\text{Ap}(S') = \{0, 79, 26, 27, 52, 53\}$$



The *Kunz poset* of S : use ground set \mathbb{Z}_m instead of $\text{Ap}(S)$.

Theorem (Bruns–García–Sánchez–O.–Wilburne)


Numerical semigroups lie in the relative interior of the same face of C_m if and only if their Kunz posets are identical.

Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

$$S = \langle 6, 9, 20 \rangle$$
$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



The *Kunz poset* of S : use ground set \mathbb{Z}_m instead of $\text{Ap}(S)$.

Theorem (Bruns–García-Sánchez–O.–Wilburne)

Numerical semigroups lie in the relative interior of the same face of C_m if and only if their Kunz posets are identical.


Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



Defining facet equations:

$$2a_2 = a_4$$

$$a_2 + a_3 = a_5$$

$$a_2 + a_5 = a_1$$

$$a_3 + a_4 = a_1$$

The *Kunz poset* of S : use ground set \mathbb{Z}_m instead of $\text{Ap}(S)$.

Theorem (Bruns–García–Sánchez–O.–Wilburne)

Numerical semigroups lie in the relative interior of the same face of C_m if and only if their Kunz posets are identical.


Faces of the Kunz cone

Question

When are numerical semigroups in (the relative interior of) the same face?

$$S = \langle 6, 9, 20 \rangle$$

$$\text{Ap}(S) = \{0, 49, 20, 9, 40, 29\}$$



Defining facet equations:


$$\begin{array}{ll} 2a_2 = a_4 & 2 \preceq 4 \\ a_2 + a_3 = a_5 & 2 \preceq 5 \\ & 3 \preceq 5 \\ a_2 + a_5 = a_1 & 2 \preceq 1 \\ & 5 \preceq 1 \\ a_3 + a_4 = a_1 & 3 \preceq 1 \\ & 4 \preceq 1 \end{array}$$

The *Kunz poset* of S : use ground set \mathbb{Z}_m instead of $\text{Ap}(S)$.


Theorem (Bruns–García–Sánchez–O.–Wilburne)

Numerical semigroups lie in the relative interior of the same face of C_m if and only if their Kunz posets are identical.

Faces of the Kunz cone




Faces of the Kunz cone



Big Q: what algebraic properties are determined by the Kunz poset P of S ?

Faces of the Kunz cone



Big Q: what algebraic properties are determined by the Kunz poset P of S ?

Spoiler

If two numerical semigroups S and S' have identical Kunz posets, then S and S' have the same number of minimal trades.

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Example

$S = \langle 6, 9, 20 \rangle$:

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Example

$S = \langle 6, 9, 20 \rangle$:

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

$$Z(1000001) =$$

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Example

$S = \langle 6, 9, 20 \rangle$:

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

$$Z(1000001) = \left\{ \underbrace{\hspace{10em}}_{\text{shortest}}, \dots, \underbrace{\hspace{10em}}_{\text{longest}} \right\}$$

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Example

$S = \langle 6, 9, 20 \rangle$:

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

$$Z(1000001) = \left\{ \underbrace{(2, 1, 49999)}_{\text{shortest}}, \dots, \underbrace{\hspace{10em}}_{\text{longest}} \right\}$$

Minimal presentations and Betti elements

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$Z(n) = \left\{ \mathbf{a} \in \mathbb{Z}_{\geq 0}^k : n = a_1 n_1 + \dots + a_k n_k \right\}$$

is the set of factorizations of $n \in S$.

Example

$S = \langle 6, 9, 20 \rangle$:

$$Z(60) = \{(10, 0, 0), (7, 2, 0), (4, 4, 0), (1, 6, 0), (0, 0, 3)\}$$

$$Z(1000001) = \left\{ \underbrace{(2, 1, 49999)}_{\text{shortest}}, \dots, \underbrace{(166662, 1, 1)}_{\text{longest}} \right\}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \cdots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\begin{aligned} \mathbf{a} &\sim \mathbf{a} \\ \mathbf{a} \sim \mathbf{b} &\Rightarrow \mathbf{b} \sim \mathbf{a} \\ \mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} &\Rightarrow \mathbf{a} \sim \mathbf{c} \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

Monomial map:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b})$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a} \quad x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$\begin{aligned} x^{\mathbf{a}} - x^{\mathbf{a}} &= 0 \in I_S \\ x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S &\Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S$$

$$(x^{\mathbf{a}} - x^{\mathbf{b}}) + (x^{\mathbf{b}} - x^{\mathbf{c}}) = x^{\mathbf{a}} - x^{\mathbf{c}}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Definition

The *kernel* $\ker \pi$ is the relation \sim on $\mathbb{Z}_{\geq 0}^k$ with $\mathbf{a} \sim \mathbf{b}$ whenever

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) \quad x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S = \ker \varphi$$

$\ker \pi$ is a *congruence*: an equivalence relation

$$\mathbf{a} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{b} \sim \mathbf{a}$$

$$\mathbf{a} \sim \mathbf{b} \text{ and } \mathbf{b} \sim \mathbf{c} \Rightarrow \mathbf{a} \sim \mathbf{c}$$

that is closed under *translation*

$$\mathbf{a} \sim \mathbf{b} \Rightarrow \mathbf{a} + \mathbf{c} \sim \mathbf{b} + \mathbf{c}$$

$$x^{\mathbf{a}} - x^{\mathbf{a}} = 0 \in I_S$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{b}} - x^{\mathbf{a}} \in I_S$$

$$(x^{\mathbf{a}} - x^{\mathbf{b}}) + (x^{\mathbf{b}} - x^{\mathbf{c}}) = x^{\mathbf{a}} - x^{\mathbf{c}}$$

$$x^{\mathbf{a}} - x^{\mathbf{b}} \in I_S \Rightarrow x^{\mathbf{c}}(x^{\mathbf{a}} - x^{\mathbf{b}}) \in I_S$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \Leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \Leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \Leftrightarrow *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle:$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$

$Z(18)$:

(3, 0, 0)



(0, 2, 0)

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$




Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$

$Z(18):$ $(3, 0, 0)$  $(0, 2, 0)$	$Z(60):$  $(10, 0, 0)$ $(7, 2, 0)$  $(0, 0, 3)$ $(4, 4, 0)$ $(1, 6, 0)$	$x^7 y^2 - x^4 y^4 = x^4 y^2 (x^3 - y^2)$ $x^7 y^2 - z^3 = (x^7 y^2 - x^4 y^4) + (x^4 y^4 - z^3)$
---	--	--

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \Leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \Leftrightarrow *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \Leftrightarrow \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \Leftrightarrow *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$

minimal presentation of S


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal generating set of I_S

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of I_S :

$$\begin{aligned} I_S &= \langle x^3 - y^2, x^{10} - z^3 \rangle \\ &= \langle x^3 - y^2, x^7 y^2 - z^3 \rangle \\ &= \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \\ &= \langle x^3 - y^2, x^6 y - z^3 \rangle \end{aligned}$$

Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



Minimal presentations

Fix a numerical semigroup $S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0}$.

$$n = a_1 n_1 + \dots + a_k n_k \quad \iff \quad \mathbf{a} = (a_1, \dots, a_k) \in \mathbb{Z}_{\geq 0}^k$$

Factorization homomorphism:

$$\begin{aligned} \pi : \mathbb{Z}_{\geq 0}^k &\longrightarrow \langle n_1, \dots, n_k \rangle \\ \mathbf{a} &\longmapsto a_1 n_1 + \dots + a_k n_k \end{aligned}$$


Monomial map:

$$\begin{aligned} \varphi : \mathbb{k}[x_1, \dots, x_k] &\longrightarrow \mathbb{k}[w] \\ x_i &\longmapsto w^{n_i} \end{aligned}$$

minimal presentation of S \iff *minimal generating set of I_S*

Example

$$S = \langle 6, 9, 20 \rangle: \quad I_S = \langle x^3 - y^2, x^4 y^4 - z^3 \rangle \subseteq \mathbb{k}[x, y, z]$$



All minimal generating sets of \sim :

$$(3, 0, 0) \sim (0, 2, 0), (10, 0, 0) \sim (0, 0, 3)$$

$$(3, 0, 0) \sim (0, 2, 0), (7, 2, 0) \sim (0, 0, 3)$$

$$(3, 0, 0) \sim (0, 2, 0), (4, 4, 0) \sim (0, 0, 3)$$

$$(3, 0, 0) \sim (0, 2, 0), (1, 6, 0) \sim (0, 0, 3)$$

Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$

Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$

Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)



Z(226)



Z(208)




Z(360)

Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)



Z(208)



Z(360)





Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$





Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)



Z(208)



Z(360)





Z(550)

Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)



Z(208)



Z(360)





Z(550)

Minimal presentations and Betti elements

$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$

$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




$\mathbb{Z}(132)$




$\mathbb{Z}(318)$




$\mathbb{Z}(226)$




$\mathbb{Z}(208)$



$\mathbb{Z}(360)$



$\mathbb{Z}(550)$




$\mathbb{Z}(800)$

Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)




Z(208)



Z(360)



Z(550)




Z(800)

Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)




Z(208)



Z(360)



Z(550)




Z(800)

Minimal presentations and Betti elements


$$S = \langle n_1, \dots, n_k \rangle \subseteq \mathbb{Z}_{\geq 0} \quad \pi : \mathbb{Z}_{\geq 0}^k \longrightarrow S$$

A larger example: $S = \langle 13, 44, 106, 120 \rangle$


$$I_S = \langle x_1^6 x_4^2 - x_3^3, x_1^2 x_3 - x_2^3, x_1^{14} x_2 - x_3 x_4, x_1^{16} - x_2^2 x_4, x_1^6 x_2^4 x_3 - x_4^3 \rangle$$




Z(132)




Z(318)




Z(226)




Z(208)



Z(360)



Z(550)



Z(800)

Minimal trades and Kunz posets


Question

How can one recover minimal trade structure from the Kunz poset?

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?




Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$




Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$


$$S = \langle 9, a_2, a_3, a_8 \rangle$$



Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?



$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$


$$S = \langle 9, a_2, a_3, a_8 \rangle$$

Cover relations: add a generator

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?



$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$

$$S = \langle 9, a_2, a_3, a_8 \rangle$$


Cover relations: add a generator

$$Z(a_6) = \{(0, 3, 0, 0), (0, 0, 2, 0)\}$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?



$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$

$$S = \langle 9, a_2, a_3, a_8 \rangle$$

Cover relations: add a generator

$$Z(a_6) = \{(0, 3, 0, 0), (0, 0, 2, 0)\}$$

2 “inner” minimal trades:


$$(0, 3, 0, 0) \sim (0, 0, 2, 0) \text{ (at } a_6)$$

$$(0, 2, 1, 0) \sim (0, 0, 0, 2) \text{ (at } a_7)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?



$$\text{Ap}(S) = \{0, a_1, a_2, \dots, a_8\}$$

$$S = \langle 9, a_2, a_3, a_8 \rangle$$

Cover relations: add a generator

$$Z(a_6) = \{(0, 3, 0, 0), (0, 0, 2, 0)\}$$

2 “inner” minimal trades:

$$(0, 3, 0, 0) \sim (0, 0, 2, 0) \text{ (at } a_6)$$

$$(0, 2, 1, 0) \sim (0, 0, 0, 2) \text{ (at } a_7)$$

Moral: can recover

- factorizations of $a \in \text{Ap}(S)$
- (minimal) trades at $a \in \text{Ap}(S)$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j


Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j

$$S = \langle 17, a_3, a_{15} \rangle$$



Minimal trades and Kunz posets


Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j

$$S = \langle 17, a_3, a_{15} \rangle$$

3 minimal trades, none in $\text{Ap}(S)$




Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 17, a_3, a_{15} \rangle$$

3 minimal trades, none in $\text{Ap}(S)$

$$a_{12} + a_3: (0, 5, 0) \sim (, ,)$$

$$a_{11} + a_{15}: (0, 0, 4) \sim (, ,)$$


$$a_2 + a_{15}: (0, 2, 3) \sim (, ,)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 17, a_3, a_{15} \rangle$$

3 minimal trades, none in $\text{Ap}(S)$

$$a_{12} + a_3: (0, 5, 0) \sim (*, 0, 1)$$

$$a_{11} + a_{15}: (0, 0, 4) \sim (*, 3, 0)$$


$$a_2 + a_{15}: (0, 2, 3) \sim (*, 0, 0)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 17, a_3, a_{15} \rangle$$

3 minimal trades, none in $\text{Ap}(S)$

$$a_{12} + a_3: (0, 5, 0) \sim (*, 0, 1)$$

$$a_{11} + a_{15}: (0, 0, 4) \sim (*, 3, 0)$$

$$a_2 + a_{15}: (0, 2, 3) \sim (*, 0, 0)$$

Possible method to locate the “outer” trades:

- factorizations of $a \in \text{Ap}(S)$ form a monomial staircase
- one “outer” minimal trade for each monomial generator

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j


Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j

$$S = \langle 10, a_2, a_3, a_4 \rangle$$




Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :


$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Candidates for “outer” trades:

$$(0, 0, 2, 1), (0, 1, 0, 2),$$


$$(0, 0, 0, 3), (0, 2, 0, 0)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Candidates for “outer” trades:

$$(0, 0, 2, 1), (0, 1, 0, 2),$$


$$(0, 0, 0, 3), (0, 2, 0, 0)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Candidates for “outer” trades:

$$(0, 0, 2, 1), (0, 1, 0, 2),$$


$$(0, 0, 0, 3), (0, 2, 0, 0)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Candidates for “outer” trades:

$$(0, 0, 2, 1), (0, 1, 0, 2),$$

$$(0, 0, 0, 3), (0, 2, 0, 0)$$


Moral: use **sets** of factorizations,
avoids overcounting minimal trades

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 10, a_2, a_3, a_4 \rangle$$

“inner” trade at a_6 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

Candidates for “outer” trades:

$$(0, 0, 2, 1), (0, 1, 0, 2),$$

$$(0, 0, 0, 3), (0, 2, 0, 0)$$

Moral: use **sets** of factorizations,
avoids overcounting minimal trades

$$0: \{(0, 0, 2, 1), (0, 1, 0, 2)\}$$

$$2: \{(0, 0, 0, 3)\}, \quad 4: \{(0, 2, 0, 0)\}$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j


Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j

$$S = \langle 6, 7, 8, 9 \rangle$$



Minimal trades and Kunz posets

Question


How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j

$$S = \langle 6, 7, 8, 9 \rangle$$

“inner” trade at a_4 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$




Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 6, 7, 8, 9 \rangle$$

“inner” trade at a_4 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

candidate for “outer” trade:


$$(0, 0, 2, 1) \in Z(25)$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 6, 7, 8, 9 \rangle$$

“inner” trade at a_4 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

candidate for “outer” trade:

$$(0, 0, 2, 1) \in Z(25)$$

No trades in $Z(25)$:


$$\{(0, 0, 2, 1), (0, 1, 0, 2), (3, 1, 0, 0)\}$$

Minimal trades and Kunz posets

Question

How can one recover minimal trade structure from the Kunz poset?

Key fact: each trade occurs at $a_i + n_j$ for some $a_i \in \text{Ap}(S)$, generator n_j



$$S = \langle 6, 7, 8, 9 \rangle$$

“inner” trade at a_4 :

$$(0, 0, 2, 0) \sim (0, 1, 0, 1)$$

candidate for “outer” trade:

$$(0, 0, 2, 1) \in Z(25)$$

No trades in $Z(25)$:

$$\{(0, 0, 2, 1), (0, 1, 0, 2), (3, 1, 0, 0)\}$$

A technical definition

Definition


An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

A technical definition

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

$$S = \langle 10, a_2, a_3, a_4 \rangle$$



$$B = \{(0, 0, 2, 1), (0, 1, 0, 2)\}$$


$$B - e_2 = \{(0, 0, 0, 2)\} = Z(a_8)$$

$$B - e_3 = \{(0, 0, 1, 1)\} = Z(a_7)$$

$$B - e_4 = \{(0, 0, 2, 0), (0, 1, 0, 1)\}$$

$$= Z(a_6)$$

$$S = \langle 6, 7, 8, 9 \rangle$$



$$B = \{(0, 0, 2, 1)\}?$$

$$B - e_4 = \{(0, 0, 2, 0)\} \subsetneq Z(a_4)$$

$$B = \{(0, 0, 2, 1), (0, 1, 0, 2)\}?$$

$$B - e_3 = \{(0, 0, 1, 1)\} \not\subseteq Z(a_i)$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Recovering minimal presentation from the Kunz poset P of S :

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Recovering minimal presentation from the Kunz poset P of S :

- trades occurring at $a \in \text{Ap}(S)$ recovered from factorizations of $\bar{a} \in P$

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Recovering minimal presentation from the Kunz poset P of S :

- trades occurring at $a \in \text{Ap}(S)$ recovered from factorizations of $\bar{a} \in P$
- each trade occurring outside $\text{Ap}(S)$ corresponds to an outer Betti element of P

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Recovering minimal presentation from the Kunz poset P of S :

- trades occurring at $a \in \text{Ap}(S)$ recovered from factorizations of $\bar{a} \in P$
- each trade occurring outside $\text{Ap}(S)$ corresponds to an outer Betti element of P

Theorem (Gomes–O.–Torres Davila)

If S has Kunz poset P , each minimal trade of S not occurring in $\text{Ap}(S)$ contains a factorization from a distinct outer Betti element of P .

In particular, if S, S' have identical Kunz poset, then S and S' have the same number of minimal trades.

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.


Another subtlety: distinct outer Betti elements can **coincide** for some S

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Another subtlety: distinct outer Betti elements can **coincide** for some S



$$B_1 = \{(0, 0, 2, 0)\}$$


$$B_2 = \{(0, 1, 0, 1)\}$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Another subtlety: distinct outer Betti elements can **coincide** for some S



$$S = \langle 4, 9, 14, 11 \rangle$$

$$B_1 = \{(0, 0, 2, 0)\}$$


$$B_2 = \{(0, 1, 0, 1)\}$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Another subtlety: distinct outer Betti elements can **coincide** for some S



$$S = \langle 4, 9, 14, 11 \rangle$$

$$20: (0, 1, 0, 1), (5, 0, 0, 0)$$

$$28: (0, 0, 2, 0), (2, 1, 0, 0), (5, 0, 0, 0)$$

$$B_1 = \{(0, 0, 2, 0)\}$$


$$B_2 = \{(0, 1, 0, 1)\}$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Another subtlety: distinct outer Betti elements can **coincide** for some S



$$S = \langle 4, 9, 14, 11 \rangle$$

$$20: (0, 1, 0, 1), (5, 0, 0, 0)$$

$$28: (0, 0, 2, 0), (2, 1, 0, 0), (5, 0, 0, 0)$$

$$S = \langle 4, 9, 10, 11 \rangle$$

$$B_1 = \{(0, 0, 2, 0)\}$$


$$B_2 = \{(0, 1, 0, 1)\}$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Another subtlety: distinct outer Betti elements can **coincide** for some S



$$B_1 = \{(0, 0, 2, 0)\}$$

$$B_2 = \{(0, 1, 0, 1)\}$$

$$S = \langle 4, 9, 14, 11 \rangle$$

$$20: (0, 1, 0, 1), (5, 0, 0, 0)$$

$$28: (0, 0, 2, 0), (2, 1, 0, 0), (5, 0, 0, 0)$$

$$S = \langle 4, 9, 10, 11 \rangle$$

$$20: (0, 0, 2, 0), (0, 1, 0, 1), (5, 0, 0, 0)$$

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Theorem (Gomes–O.–Torres Davila)

If S has Kunz poset P , each minimal trade of S not occurring in $\text{Ap}(S)$ contains a factorization from a distinct outer Betti element of P .

In particular, if S, S' have identical Kunz poset, then S and S' have the same number of minimal trades.

The main theorem

Definition

An *outer Betti element* of a Kunz poset P is a set B of factorizations with connected factorization graph and $B - e_i = Z(a_i)$ for each $i \in \text{supp}(B)$.

Theorem (Gomes–O.–Torres Davila)

If S has Kunz poset P , each minimal trade of S not occurring in $\text{Ap}(S)$ contains a factorization from a distinct outer Betti element of P .

In particular, if S, S' have identical Kunz poset, then S and S' have the same number of minimal trades.

For $m = 6$: $\#$ minimal trades $\in \{1, 2, 3, 4, 5, 6, 9, 10, 15\}$

Application: classifying minimal trades


Question

Given the multiplicity $m = m(S)$ and $\#$ minimal generators k of a numerical semigroup S , what can $\beta_1(I_S) = \#$ minimal trades be?

Application: classifying minimal trades

Question

Given the multiplicity $m = m(S)$ and $\#$ minimal generators k of a numerical semigroup S , what can $\beta_1(I_S) = \#$ minimal trades be?




Prior work: a family has $\beta_1(S) = \binom{k}{2}$ for $3 \leq k \leq m$ (Rosales)

if $r = m - k \leq 2$, then $\beta_1(S) \in [\binom{k}{2} - r, \binom{k}{2}]$ (GS-R)

Application: classifying minimal trades

Question

Given the multiplicity $m = m(S)$ and $\#$ minimal generators k of a numerical semigroup S , what can $\beta_1(I_S) = \#$ minimal trades be?



Using Kunz posets: a family hits each $\beta_1(S) \in [\binom{k}{2} - r, \binom{k}{2}]$


for $r = m - k \leq k - 2$

a family hits $\beta_1(S) = \binom{k}{2} + 1$ for each $m \geq k + 3$

Application: classifying minimal trades

Question

Given the multiplicity $m = m(S)$ and $\#$ minimal generators k of a numerical semigroup S , what can $\beta_1(I_S) = \#$ minimal trades be?







Bounds from Kunz posets: $\beta_1(S) \geq \binom{k}{2} - r$, where $r = m - k$
 if $m - k = 3$, then $\beta_1(S) \in [\binom{k}{2} - 3, \binom{k}{2} + 1]$

References

-  W. Bruns, P. García-Sánchez, C. O'Neill, D. Wilburne (2020)
Wilf's conjecture in fixed multiplicity
International Journal of Algebra and Computation **30** (2020), no. 4, 861–882.
(arXiv:1903.04342)
-  N. Kaplan, C. O'Neill, (2021)
Numerical semigroups, polyhedra, and posets I: the group cone
Combinatorial Theory **1** (2021), #19. (arXiv:1912.03741)
-  T. Gomes, C. O'Neill, E. Torres Davila (2022)
Numerical semigroups, polyhedra, and posets III: minimal presentations and face dimension.
Electronic Journal of Combinatorics **30** (2023), no. 2, #P2.5. (arXiv:2009.05921)
-  C. Elmacioglu, K. Hilmer, H. Koufmann, C. O'Neill, M. Okandan (2022)
On the cardinality of minimal presentations of numerical semigroups
under review. (arXiv:2211.16283)

References

-  W. Bruns, P. García-Sánchez, C. O'Neill, D. Wilburne (2020)
Wilf's conjecture in fixed multiplicity
International Journal of Algebra and Computation **30** (2020), no. 4, 861–882.
(arXiv:1903.04342)
-  N. Kaplan, C. O'Neill, (2021)
Numerical semigroups, polyhedra, and posets I: the group cone
Combinatorial Theory **1** (2021), #19. (arXiv:1912.03741)
-  T. Gomes, C. O'Neill, E. Torres Davila (2022)
Numerical semigroups, polyhedra, and posets III: minimal presentations and face dimension.
Electronic Journal of Combinatorics **30** (2023), no. 2, #P2.5. (arXiv:2009.05921)
-  C. Elmacioglu, K. Hilmer, H. Koufmann, C. O'Neill, M. Okandan (2022)
On the cardinality of minimal presentations of numerical semigroups
under review. (arXiv:2211.16283)

Thanks!