

Fall 2020, Math 579: Problem Set 7
Due: Thursday, October 15th, 2020
Partitions

Discussion problems. The problems below should be worked on in class.

(D1) *Bell numbers.* Recall that $S(n, k)$ denotes the number of set partitions of $[n]$ into exactly k blocks (called the *Stirling numbers of the 2nd kind*). Let $B(n)$ denote the total number of set partitions of $[n]$, that is, $B(n) = \sum_{k=1}^n S(n, k)$ (called the n -th *Bell number*).

- (a) Find $B(1)$, $B(2)$, $B(3)$, and $B(4)$. Which of these did you find in the prelim problem?
- (b) Find $B(5)$ using the identity

$$B(n+1) = 1 + \sum_{k=1}^n \binom{n}{k} B(k).$$

- (c) Give a combinatorial proof that for all $n \geq 1$,

$$B(n+1) - B(n) = \sum_{k=1}^n kS(n, k).$$

(D2) *An unfortunate formula for Stirling numbers.* For $n \geq k \geq 1$, let $O_{n,k}$ denote the total number of surjective functions $[n] \rightarrow [k]$ (that is, functions in which every element of $[k]$ is the image of some element of $[n]$).

- (a) Find **all** functions $[3] \rightarrow [2]$. Label each as “surjective” or “not surjective” accordingly.
- (b) Find a formula for the total number of functions $[n] \rightarrow [k]$.
- (c) Show that $O_{5,2} = 5^2 - 2$. Hint: why is it expressed like this?
- (d) Argue that $O_{4,3} = 4^3 - \binom{3}{2}4^2 + \binom{3}{1}4^1$.
- (e) Using the idea in the previous part, find an expression for $O_{5,3}$.
- (f) For $n \geq k$, use the Sieve formula to show that

$$O_{n,k} = \sum_{j=0}^k (-1)^j \binom{k}{j} (k-j)^n.$$

Hint: first, verify this expression matches what you found in the previous part.

- (g) Verify $O_{4,3} = \binom{4}{2} \cdot 3 \cdot 2$. Hint: why is it expressed like this?
- (h) Give a combinatorial proof that $S(n, k) = O_{n,k}/k!$.

Homework problems. You must submit *all* homework problems in order to receive full credit.


- (H1) Use the recurrence identity $S(n, k) = S(n-1, k-1) + k \cdot S(n-1, k)$ from class to complete the next 3 rows of the triangle of Stirling numbers of the 2nd kind (the analog of Pascal's triangle where the entry in row n and position k is $S(n, k)$).

$$\begin{array}{ccccccc}
 & & & & 1 & & & & \\
 & & & & 1 & & 1 & & \\
 & & & 1 & & 3 & & 1 & \\
 & & 1 & & 7 & & 6 & & 1
 \end{array}$$

- (H2) Find a closed formula for $S(n, n-2)$ in terms of n , valid for $n \geq 3$.
- (H3) Find a formula for the number of partitions of $[n]$ into blocks of size exactly 2.
- (H4) Let $F(n)$ denote the number of set partitions of $[n]$ with no singleton blocks. Prove that $B(n) = F(n) + F(n+1)$.
- (H5) Find a recursive formula for $F(n+1)$ in terms of $F(k)$ for $k \leq n$, and give a combinatorial proof of its correctness.

Challenge problems. Challenge problems are not required for submission, but bonus points will be awarded for submitting a partial attempt or a complete solution.

- (C1) Consider the sequence of tiled hexagons below, with side lengths $n = 1, 2, 3, 4$, respectively.



Let h_n and r_n denote the number of hexagons and rhombi in the n -th tiling, respectively. Find h_n and r_n in terms of n .

Note: this problem is brought to you by Michael O'Sullivan (our department chair) and his grandson's tile set.