

Fall 2020, Math 620: Week 10 Problem Set
Due: Thursday, November 5th, 2020
A Hierarchy of Integral Domains

Discussion problems. The problems below should be completed in class.

(D1) *Euclidean domains.* In this problem, we introduce Euclidean domains.

- (a) Let $R = \mathbb{Z}$. For each a, b below, find $q, r \in R$ so that $a = qb + r$ with $0 \leq r < b$.
- (i) $a = 17, b = 3$. (ii) $a = 15, b = 5$. (iii) $a = -17, b = 5$.
- (b) Let $R = \mathbb{Q}[x]$. For each a, b below, find $q, r \in R$ so that $a = qb + r$ with $\deg(r) < \deg(b)$.
- (i) $a = x^5 + 3x^4 + 4x + 1,$ (ii) $a = x^3 + 3x^2 + 2x + 1,$
 $b = x^2 + 2x + 3.$ $b = 2x^2 + x + 3.$
- (c) Let $R = \mathbb{Z}_5[x]$. For each a, b below, find $q, r \in R$ so that $a = qb + r$ with $\deg(r) < \deg(b)$.
- (i) $a = x^5 + 3x^4 + 4x + 1,$ (ii) $a = x^3 + 3x^2 + 2x + 1,$
 $b = x^2 + 2x + 3.$ $b = 2x^2 + x + 3.$
- (d) Will the division algorithm work in $F[x]$ for any field F ? Briefly justify your answer.
- (e) Let $R = \mathbb{Z}[i]$. For each a, b below, find $q, r \in R$ so that $a = qb + r$ with $\|r\| < \|b\|$.
- (i) $a = 1 + 21i,$ (ii) $a = 10 + 15i,$ (iii) $a = 2 + 23i,$
 $b = 2 + 3i.$ $b = 4 + 6i.$ $b = 1 + 2i.$

Hint: for part (i), a remainder of 0 is possible.

- (f) Are your remainders in part (e) unique?
- (g) A *Euclidean domain* is an integral domain R equipped with a *norm* $N : R \setminus \{0\} \rightarrow \mathbb{Z}_{\geq 0}$ such that for every $a, b \in R$ with $b \neq 0$, there exists $q, r \in R$ with $r = 0$ or $N(r) < N(b)$ so that $a = qb + r$. Identify the norm function of each ring above.

(D2) *Greatest common divisors.* The goal of this problem is explore $\gcd()$ for Euclidean domains.

- (a) Let $R = \mathbb{Z}$. Find $\gcd(42, 96)$ using the Euclidean algorithm.
- (b) Show that the ideal $\langle 42, 96 \rangle \subset \mathbb{Z}$ is principle.
- (c) Let $R = \mathbb{Z}_3[x]$. Find $\gcd(x^6 + x^4 + x^2, x^4 + x^3 + x)$ using the Euclidean algorithm.
- (d) Show that the ideal $\langle x^6 + x^4 + x^2, x^4 + x^3 + x \rangle \subset \mathbb{Z}_3[x]$ is principle.
- (e) Propose a definition for **a** (not **the**) “greatest common divisor” of $a, b \in R$ for any **integral domain** R .
- (f) Argue that any Euclidean domain is a PID.

Homework problems. You must submit *all* homework problems in order to receive full credit.

(H1) Fix $D \in \mathbb{Z}_{>0}$, and let $R = \mathbb{Z}[\sqrt{-D}]$. Consider the function $N : R \setminus \{0\} \rightarrow \mathbb{Z}$ given by

$$N(a + b\sqrt{-D}) = a^2 + Db^2$$

for $a, b \in \mathbb{Z}$.

- (a) Prove that $N(zw) = N(z)N(w)$ for any $z, w \in R$.
 - (b) Prove that $z \in R$ is a unit if and only if $N(z) = 1$.
 - (c) Prove that if $D = 5$, then R is not a UFD.
 - (d) For $D = 2$, determine if R is a Euclidean domain, a PID, a UFD, or none of these.
- (H2) Suppose F is a field, and fix $f(x) \in F[x]$ and $a \in F$. Prove that $f(a) = 0$ if and only if $f(x) = (x - a)g(x)$ for some $g(x) \in F[x]$.

(H3) Consider the ring

$$R = \{f(x) \in \mathbb{Q}[x] : f(n) \in \mathbb{Z} \text{ for all } n \in \mathbb{Z}\}$$

of *integer valued polynomials*.

- (a) Prove that R is a ring with $\mathbb{Z}[x] \subsetneq R \subsetneq \mathbb{Q}[x]$.
- (b) Prove that R is not a UFD.

(H4) Consider the integral domain

$$R = \{f(x) \in \mathbb{Q}[x] : f(0) \in \mathbb{Z}\}$$

consisting of polynomials with rational coefficients and integer constant term. Prove that the polynomial $x \in R$ cannot be written as a product of finitely many irreducible elements of R (we say R is not *atomic*).

Challenge problems. Challenge problems are not required for submission, but bonus points will be awarded for submitting a partial attempt or a complete solution.

(C1) Prove or disprove: if $I \subset \mathbb{Z}[i]$ is any nontrivial ideal, then $\mathbb{Z}[i]/I$ has finitely many elements.