

**Fall 2022, Math 522: Week 5 Problem Set**  
**Due: Friday, October 7th, 2022**  
**Modular Arithmetic (Week 2)**

**Discussion problems.** The problems below should be worked on in class.

- (D1) *The orders of elements of  $\mathbb{Z}_n$ .* The *order* of an element  $[a]_n \in \mathbb{Z}_n$  is the smallest integer  $k$  such that adding  $[a]_n$  to itself  $k$  times yields  $[0]_n$ , that is,  $ka \equiv 0 \pmod n$ .
- (a) Find the order of each element of  $\mathbb{Z}_{12}$ . Do the same for  $\mathbb{Z}_{10}$ .
  - (b) Conjecture a formula for the order of  $[a]_n$  in terms of  $a$  and  $n$ .  
Hint: use your answers from part (a) for inspiration. When in doubt, do more examples!
  - (c) Let  $k$  denote your conjectured order for  $[a]_n$ . Prove  $[k]_n[a]_n = 0$ .
  - (d) Let  $k$  denote your conjectured order for  $[a]_n$ , and suppose  $[c]_n[a]_n = 0$ . Prove  $k \mid c$ .  
Hint: how have we used the division algorithm to prove one integer divides another?
  - (e) Prove that your conjectured order formula holds.
  - (f) For which  $n$  does every nonzero  $[a]_n$  have order  $n$ ? Give a (short and sweet) proof.
- (D2) *Euler's theorem.* Fix  $n \geq 1$ , and let  $s = \phi(n)$  denote the number of integers  $i \in [1, n - 1]$  with  $\gcd(i, n) = 1$  (this is known as the *Euler totient function*). The goal of this problem is to prove the following theorem.

**Theorem** (Euler's Theorem). *If  $\gcd(a, n) = 1$ , then  $a^s \equiv 1 \pmod n$ .*

- (a) Compare your answers to the prelim problem. Write the definition of *reduced residue system* at the top of the board, along with at least two examples.
- (b) Prove that if  $r_1, \dots, r_s$  is some reduced residue system for  $n$  and  $\gcd(a, n) = 1$ , then  $ar_1, \dots, ar_s$  is also a reduced residue system for  $n$ .  
Hint: the "cancellation law" should come in handy somewhere in your proof.
- (c) What does part (b) tell you about the products  $r_1 \cdots r_s$  and  $(ar_1) \cdots (ar_s)$  modulo  $n$ ?
- (d) Conclude that Euler's theorem holds.
- (e) Use Euler's theorem to prove Fermat's little theorem.

**Homework problems.** You must submit *all* homework problems in order to receive full credit.

Unless otherwise stated,  $a, b, c, n, p \in \mathbb{Z}$  are arbitrary with  $p > 1$  prime and  $n \geq 2$ .

- (H1) Determine how many primes  $p$  satisfy  $n! + 2 \leq p \leq n! + n$ . Prove your claim.
- (H2) Prove that  $10 \nmid (n-1)! + 1$  for all  $n \geq 1$ . What does this tell you about the hypotheses for Wilson's theorem?
- (H3) Prove that if  $\gcd(a, n) = \gcd(a-1, n) = 1$ , then  $1 + a + a^2 + \cdots + a^{\phi(n)-1} \equiv 0 \pmod{n}$ .
- (H4) Prove that if  $p$  is prime, then  $(a+b)^p \equiv a^p + b^p \pmod{p}$  for every  $a, b \in \mathbb{Z}$  (this is known as the *Freshmen's Dream*).

Note: you may **not** use the binomial theorem in this problem.

- (H5) Determine whether each of the following is true or false. Prove each true statement, and give a counterexample for each false statement.
  - (a) If  $\gcd(a, n) = 1$  and  $a \not\equiv 1 \pmod{n}$ , then the smallest positive integer  $b$  such that  $a^b \equiv 1 \pmod{n}$  is  $b = \phi(n)$ .
  - (b) If  $n \geq 2$ , then  $(a+b)^n \equiv a^n + b^n \pmod{n}$  for every  $a, b \in \mathbb{Z}$ .