

Spring 2019, Math 320: Week 11 Problem Set
Due: Tuesday, April 23rd, 2019
Congruence Classes in $F[x]$ (Week 1)

Discussion problems. The problems below should be worked on in class.

(D1) *Arithmetic modulo $p(x)$.* For this problem, all polynomials have coefficients in \mathbb{Q} .

- (a) Determine whether $x^3 + 2x + 1 \equiv x^2 + 1 \pmod{x^2 - 1}$.
- (b) Locate the canonical representative of $x^3 + 2x + 1$ modulo $x^2 - 1$.
- (c) Use the fact that $[x^2 - 1] = [0]$ in $\mathbb{Q}[x]/\langle x^2 - 1 \rangle$ to demonstrate $[x]^2 = [1]$.
- (d) Use part (c) and congruence class arithmetic to determine if $[x^3 + 2x + 1] = [x^2 + 1]$.
- (e) Demonstrate $[x - 1] \in \mathbb{Q}[x]/\langle x^2 - 2x + 1 \rangle$ is a zero-divisor.
- (f) Demonstrate $[x - 2] \in \mathbb{Q}[x]/\langle x^2 - 2x + 1 \rangle$ is a unit.

(D2) *Interpreting quotient rings.* Consider the rings

$$R = \mathbb{Q}[\sqrt{2}] = \{a + b\sqrt{2} : a, b \in \mathbb{Q}\} \subset \mathbb{R} \quad \text{and} \quad T = \mathbb{Q}[x]/\langle x^2 - 2 \rangle.$$

- (a) Determine whether $\mathbb{Q} \subset R$. Determine whether $\sqrt{3} \in R$.
- (b) Find the sum and product of $2 + 3\sqrt{2}$ and $5 + 7\sqrt{2}$ in R .
- (c) Find the canonical representative of $[x^2]$ in T .
- (d) Find the sum and product of $[2 + 3x]$ and $[5 + 7x]$ in T . Do they look familiar?
- (e) Define $\varphi : R \rightarrow T$ by $\varphi(a + b\sqrt{2}) = [a + bx]$. We will show φ is an isomorphism.
 - (i) Begin by proving φ is a homomorphism.
 - (ii) Fill in the gaps in the following proof that φ is injective.

Proof. Suppose $\varphi(a + b\sqrt{2}) = \varphi(c + d\sqrt{2})$. This implies $[a + bx] = [c + dx]$, so

$$(a - c) + (b - d)x = (a + bx) - (c + dx) = \underline{\hspace{2cm}}$$

for some $k(x) \in \mathbb{Q}[x]$. Since $\underline{\hspace{2cm}}$, we must have $k(x) = 0$.
This yields $a - c = 0$ and $b - d = 0$, and thus $a + b\sqrt{2} = c + d\sqrt{2}$. □

- (iii) Lastly, complete the proof that φ is an isomorphism.

(D3) *Identifying familiar quotient rings.*

- (a) Find a field F and a polynomial $p(x)$ so that $\mathbb{Q}[\sqrt{5}] \cong F[x]/\langle p(x) \rangle$.
- (b) Consider the set

$$R = \{a + b\sqrt[3]{2} : a, b \in \mathbb{Q}\} \subset \mathbb{R}.$$

Is R a ring? If not, what must be added to ensure R is closed under both operations?

- (c) Based on part (b), how should $\mathbb{Q}[\sqrt[3]{2}]$ be defined so that it is a ring?
- (d) Find a field F and a polynomial $p(x)$ so that $\mathbb{Q}[\sqrt[3]{2}] \cong F[x]/\langle p(x) \rangle$. No proof is required, but specify the isomorphism you would use.
- (e) Fix a field F , $a \in F$, and $f(x) \in F[x]$, and let $R = F[x]/\langle x - a \rangle$. Locate and correct the (subtle!) error in the proof that the canonical representative of $[f(x)]$ is $[f(a)]$.

Proof. Write $f(x) = b_d x^d + \cdots + b_1 x + b_0$, where $d \in \mathbb{Z}_{\geq 0}$ and $b_0, \dots, b_d \in F$. Since $[x - a] = [0]$, we have $[x] = [a]$, so

$$\begin{aligned} [f(x)] &= [b_d x^d + \cdots + b_1 x + b_0] = [b_d][x]^d + \cdots + [b_1][x] + [b_0] \\ &= [b_d][a]^d + \cdots + [b_1][a] + [b_0] = [b_d a^d + \cdots + b_1 a + b_0] = [f(a)] \end{aligned}$$

Since $f(a)$ is a constant, $\deg f(a) = 0$, so $\deg f(a) < \deg(x - a)$, as desired. □

Homework problems. As the name suggests, you must submit *all* required problem with this homework set in order to receive full credit.

- (H1) Determine whether $x^3 + x^2 \equiv x \pmod{(x^2 + x + 1)}$ over \mathbb{Z}_2 . Do the same over \mathbb{Z}_3 .
- (H2) Every element of $R = \mathbb{Q}[x]/\langle x^2 + 2x - 1 \rangle$ can be written in the form $[ax + b]$ for some $a, b \in \mathbb{Q}$ (this is the canonical representative). If $[ax + b][cx + d] = [rx + t]$ with $a, b, c, d, r, t \in \mathbb{Q}$, find formulas for r and t in terms of a, b, c , and d (this is the *multiplication rule* for R).
- (H3) Prove the map $\varphi : \mathbb{C} \rightarrow \mathbb{R}[x]/\langle x^2 + 1 \rangle$ given by $a + bi \mapsto [a + bx]$ is an isomorphism.
- (H4) Determine whether each of the following statements is true or false. Prove each true statement, and give a counterexample for each false statement.
- (a) The element $[x - 2] \in \mathbb{Q}[x]/\langle x^2 - 2x \rangle$ is a zero-divisor.
 - (b) The element $[x - 2] \in \mathbb{Q}[x]/\langle x^2 - 2x \rangle$ is a unit.
 - (c) If F is a field, $p(x) \in F[x]$, and $R = F[x]/\langle p(x) \rangle$, then R is a field.

Challenge problems. Challenge problems are not required for submission, but bonus points will be awarded for submitting a partial attempt or a complete solution.

- (C1) Consider the quotient ring $R = \mathbb{Q}[x]/\langle x^2 \rangle$. It turns out every nonzero element of R is either a unit or a zero-divisor. Determine which elements are units and which are zero-divisors.