Spring 2022, Math 579: Week 6 Problem Set Due: Thursday, March 10th, 2022 Set Partitions

Discussion problems. The problems below should be worked on in class.

- (D1) A Pascal-like recurrence. Recall that S(n,k) denotes the number of set partitions of [n] into exactly k blocks (called the Stirling numbers of the 2nd kind).
 - (a) Find S(5,4), S(4,3), and S(4,4) by listing set partitions.
 - (b) Verify that S(n+1,k) = S(n,k-1) + kS(n,k) holds for n and k in the previous part.
 - (c) Give a combinatorial proof that

$$S(n+1,k) = S(n,k-1) + kS(n,k)$$

holds for $n \ge k \ge 1$.

(D2) Bell numbers. Let B(n) denote the total number of set partitions of [n], that is,

$$B(n) = \sum_{k=1}^{n} S(n, k)$$

(called the *n*-th *Bell number*).

- (a) Find B(1), B(2), B(3), and B(4). Which of these did you find in the prelim problem?
- (b) Find B(5) using the identity

$$B(n+1) = 1 + \sum_{k=1}^{n} {n \choose k} B(k).$$

- (c) Give a combinatorial proof of the identity in part (b).
- (d) Give a combinatorial proof that for all $n \geq 1$,

$$B(n+1) - B(n) = \sum_{k=1}^{n} kS(n,k).$$

- (D3) An unfortunate formula for Stirling numbers. For $n \geq k \geq 1$, let $O_{n,k}$ denote the total number of surjective functions $[n] \rightarrow [k]$ (that is, functions in which every element of [k] is the image of some element of [n]).
 - (a) Find all functions [3] \rightarrow [2]. Label each as "surjective" or "not surjective" accordingly.
 - (b) Find a formula for the total number of functions $[n] \to [k]$.
 - (c) Show that $O_{5,2} = 5^2 2$. Hint: why is it expressed like this?
 - (d) Argue that $O_{4,3} = 4^3 {3 \choose 2} 4^2 + {3 \choose 1} 4^1$.
 - (e) Using the idea in the previous part, find an expression for $O_{5,3}$.
 - (f) For $n \geq k$, use the Sieve formula to show that

$$O_{n,k} = \sum_{j=0}^{k} (-1)^j \binom{k}{j} (k-j)^n.$$

Hint: first, verify this expression matches hat you found in the previous part.

- (g) Verify $O_{4,3} = \binom{4}{2} \cdot 3 \cdot 2$. Hint: why is it expressed like this?
- (h) Give a combinatorial proof that $S(n,k) = O_{n,k}/k!$.

Homework problems. You must submit *all* homework problems in order to receive full credit.

(H1) Use the recurrence identity $S(n,k) = S(n-1,k-1) + k \cdot S(n-1,k)$ from class to complete the next 3 rows of the triangle of Stirling numbers of the 2nd kind (the analog of Pascal's triangle where the entry in row n and position k is S(n,k)).




- (H2) Find a closed formula for S(n, n-2) in terms of n, valid for $n \geq 3$.
- (H3) Find a formula for the number of partitions of [n] into blocks of size exactly 2.
- (H4) Let F(n) denote the number of set partitions of [n] with no singleton blocks. Prove that B(n) = F(n) + F(n+1).
- (H5) Find a recursive formula for F(n+1) in terms of F(k) for $k \le n$, and give a combinatorial proof of its correctness.

Challenge problems. Challenge problems are not required for submission, but bonus points will be awarded for submitting a partial attempt or a complete solution.

(C1) Give a combinatorial proof that

$$B(n+m) = \sum_{k=1}^{n} \sum_{j=1}^{m} S(m,j) \binom{n}{k} j^{n-k} B(k).$$

(C2) Consider the sequence of tiled hexagons below, with side lengths n = 1, 2, 3, 4, respectively.



Let h_n and r_n denote the number of hexagons and rhombi in the n-th tiling, respectively. Find formulas for h_n and r_n in terms of n.

Note: this problem is brought to you by Michael O'Sullivan (our department chair) and his grandson's tile set.

2