

Spring 2026, Math 590: Week 3 Problem Set
Due: Wednesday, February 18th, 2026
Generating Functions - Deep Cuts

Discussion problems. The problems below should be worked on in class.

(D1) *Recurrence relations.* For each of the following, (i) compute a_0, \dots, a_5 , (ii) use generating functions to find a formula for a_n in terms of n , and (iii) verify your formula for $n \leq 5$.

(a) $a_0 = 1, a_1 = 3, a_n = 3a_{n-2}$

(b) $a_0 = 3, a_1 = 1, a_n = 2a_{n-1} + 3a_{n-2}$

(D2) *Power series derivatives.* Fix formal power series $A(z)$ and $B(z)$. The goal of this problem is to prove some of the “derivative rules” from Calculus.

(a) Prove that if $k \in \mathbb{Z}_{\geq 1}$, then $\frac{d}{dz}[(B(z))^k] = k(B(z))^{k-1}B'(z)$.

Hint: this can be done without writing any sigma-sums, instead using product rule and induction on k .

(b) Prove that if $B(z)$ has constant term **zero**, then $\frac{d}{dz}[A(B(z))] = A'(B(z))B'(z)$.

Hint: use part (a) and sigma-sums, but avoid writing “ b_n ” at all costs.

(D3) *Generating functions of polynomials.* We saw in class that for each $d \geq 0$, there exists a polynomial $Q_d(z)$ such that

$$\sum_{n \geq 0} n^d z^n = \frac{Q_d(z)}{(1-z)^{d+1}}.$$

(a) Write down $Q_1(z)$, $Q_2(z)$, and $Q_3(z)$ for reference (you may consult your notes).

(b) Using formal derivatives, obtain an expression for $Q_d(z)$ in terms of $Q_{d-1}(z)$.

(c) Use your expression from part (b) to show that $Q_d(z)$ has degree d .

(d) Use your expression from part (b) to show that $Q_d(z)$ has no constant term.

(e) Find a formula for $Q_d(1)$. Use your expression from part (b) to verify this.

Homework problems. You must submit *all* homework problems in order to receive full credit.

(H1) Use generating functions to find a formula for a_n if $a_0 = 1$ and $a_n = 3a_{n-1} + 2^n$.

(H2) The goal of this problem is to prove the *quotient rule* for power series.

(a) Prove that if $B(z)$ has a **nonzero** constant term, then

$$\frac{d}{dz} [(B(z))^{-1}] = \frac{-B'(z)}{(B(z))^2}.$$

Hint: to save a LOT of algebra, take the derivative of both sides of $B(z)(B(z))^{-1} = 1$ using the product rule.

(b) Prove that if $B(z)$ has a **nonzero** constant term, then

$$\frac{d}{dz} \left[\frac{A(z)}{B(z)} \right] = \frac{A'(z)B(z) - A(z)B'(z)}{(B(z))^2}.$$

Hint: using part (a) and the product rule can save a LOT of algebra.

(H3) Resuming notation from Problem (D3)(b), recall that

$$Q_1(z) = z \quad \text{and} \quad Q_d(z) = z(1-z)Q'_{d-1}(z) + dzQ_{d-1}(z)$$

for $d \geq 2$.

(a) Prove that Q_d has degree d , and that the coefficients of the positive-degree terms of $Q_d(z)$ are positive integers.

(b) Find a formula for $Q_d(1)$, and prove that your formula holds.

Challenge problems. Challenge problems are not required for submission, but bonus points will be awarded for submitting a partial attempt or a complete solution.

(C1) Prove the coefficients of $Q_d(z)$ are symmetric (that is, if we write $Q_d(z) = a_d z^d + \cdots + a_1 z$, then $a_i = a_{d+1-i}$ for each i).